

# MiR-124 affects the apoptosis of brain vascular endothelial cells and ROS production through regulating PI3K/AKT signaling pathway

S.-W. WANG, L.-X. DENG, H.-Y. CHEN, Z.-Q. SU, S.-L. YE, W.-Y. XU

Department of Neurology, the Third Hospital of Xiamen, Xiamen, Fujian, China

**Abstract.** – **OBJECTIVE:** The apoptosis of vascular endothelial cells (VEC) is related to ischemic stroke. Phosphatidylinositol-3 kinase (PI3K)/protein kinase B (AKT/PKB) signaling pathway can upregulate Bcl-2 expression, reduce reactive oxygen species (ROS) production, and induce apoptosis. The level of miR-124 was significantly increased after cerebral ischemia. This study aimed to investigate the role of miR-124 in regulating PI3K expression, brain VEC apoptosis, and ROS production.

**MATERIALS AND METHODS:** The expressions of miR-124, PI3K, p-AKT, and Bcl-2 in brain VEC of rats from the sham group and middle cerebral artery occlusion (MCAO) group were tested. Bioinformatics analysis showed the complementary binding site between miR-124 and PI3K mRNA. ROS content and cell apoptosis were detected by flow cytometry. Rat brain VEC were cultured in vitro and treated by oxygen-glucose deprivation (OGD) for 6 h. VECs were divided into four groups, including miR-NC, miR-124 inhibitor, pIRES2-blank, and pIRES2-PI3K groups, and were further treated by OGD.

**RESULTS:** MiR-124 expression, ROS content, and cell apoptosis were markedly increased, whereas the levels of PI3K, p-AKT, and Bcl-2 were markedly reduced in rat VECs from MCAO group compared with that in the sham group. OGD treatment significantly induced VECs apoptosis, upregulated miR-124 expression and ROS content, and down-regulated the levels of PI3K, p-AKT, and Bcl-2. MiR-124 inhibitor or transfection of pIRES2-PI3K plasmid apparently enhanced PI3K, p-AKT, and Bcl-2 expressions, alleviated cell apoptosis and decreased ROS content in VECs induced by OGD.

**CONCLUSIONS:** Our data demonstrated that miR-124 induced the apoptosis of brain vascular endothelial cells via the down-regulation of PI3K/AKT signaling pathway and promotion of ROS production.

*Key Words:*

Ischemic stroke, MiR-124, PI3K/AKT, Brain vascular endothelial cell.

## Introduction

Cerebral vascular disease (CVD) is a type of important disease that presents the major cause of disability and death. Ischemic CVD (ICVD) represents a common cerebrovascular disease accounting for 70-80% of stroke. It is featured as high morbidity, mortality, and disability rates, resulting in heavy burden to family and society<sup>1,2</sup>.

Vascular endothelial cells (VECs) play a crucial role in regulating vascular inflammation, maintaining the homeostasis of the vascular system and the balance between coagulation and fibrinolytic system<sup>3,4</sup>. Ischemic stroke can induce VECs injury, which aggravates the brain damage. Therefore, it is closely related to the occurrence and development of ischemic stroke<sup>5-7</sup>. Reactive oxygen species (ROS), as an important oxidative stress molecule, participates in promoting the occurrence of ischemic stroke through the mediation of VECs damage<sup>8,9</sup>. Phosphatidylinositol-3 kinase (PI3K)/protein kinase B (AKT/PKB) signaling pathway is involved in multiple biological processes, such as cell proliferation, survival, and apoptosis<sup>10</sup>. Several studies<sup>11-13</sup> showed that PI3K/AKT signaling pathway positively affected B-cell lymphoma-2 (Bcl-2) expression, alleviated oxidative damage and cell apoptosis. MiRNA is a type of endogenous single-stranded noncoding RNA at the length of 22-25 nt from the eukaryote. It exerts to the degrading or inhibiting function by binding with the 3'-UTR of targeted RNA<sup>14</sup>. MiRNA is involved in regulating cell proliferation, survival, apoptosis, oxidative stress, and ROS production. Thus, it is related to atherosclerosis<sup>15</sup>, coronary heart disease<sup>16</sup>, stroke<sup>17</sup>, and cancer<sup>18,19</sup>. Accumulative evidence revealed that the level of miR-124 was significantly elevated after ischemic brain injury<sup>20-22</sup>. This study aimed to investigate the mechanism of miR-124 in regulating the proliferation of brain vascular endothelial cells.

## Materials and Methods

### *Main Reagents and Materials*

Healthy male SD rats at 6-8 weeks old and weighted 220-250 g were purchased from Selleck (Shanghai, China). Dulbecco's Modified Eagle Medium (DMEM) medium, Fetal bovine serum (FBS), and collagen II were bought from Gibco (Waltham, MA, USA). RNA extraction kit was obtained from Roche (Indianapolis, IN, USA). Rabbit anti-rat PI3K antibody was derived from Santa Cruz Biotechnology (Santa Cruz, CA, USA). Rabbit anti-rat Bcl-2, AKT, p-AKT, and  $\beta$ -actin antibodies were got from Abcam (Cambridge, MA, USA). Horseradish peroxidase (HRP) conjugated secondary antibody was provided by Jackson ImmunoResearch (West Grove, PA, USA). Earle's balanced salt solution was prepared by HAICR (Nanjing, China). Glucose was purchased from Sigma (Temecula, CA, USA). The pGRE-luc luciferase reporter plasmid RIPA, cell apoptosis detection kit, and ROS detection kit were bought from Beyotime (Beijing, China). Lipofectamine 2000 was obtained from Thermo Fisher Scientific (Waltham, MA, USA). QuantiTect SYBR Green RT-PCR Kit was got from Qiagen (Venlo, Netherlands). Luciferase activity detection kit was provided by Promega (Fitchburg, WI, USA). MiR-NC and miR-124 inhibitor were derived from Ribobio (Guangzhou, Guangdong, China). pIRES2 overexpression plasmid was purchased from Youbio (Changsha, Hunan, China).

Rats were used for all experiments, and all procedures were approved by the Animal Ethics Committee of the third hospital of Xiamen.

### *Rat Middle Cerebral Artery Occlusion (MCAO) Model Establishment*

The rat was anesthetized with 10% chloral hydrate and fixed in supine. The neck was incised after sterilization to expose right common carotid artery, external carotid artery, and internal carotid artery. The artery clamp was used to clip the bifurcate of internal carotid and external carotid. A small incision was made using ophthalmic scissors on the internal carotid close to the intersection. The MCAO wire was inserted through the incision into the internal carotid for about 18 mm, suggesting the wire caused right MCAO. After the obligate wire was tightened and the redundant thread was cut off, the incision was sutured and the rat was treated with antibiotics. The rat in Sham group was treated by carotid artery exposed wi-

thout MCAO wire. The rat was sacrificed after three days and brain tissue and blood vessel were collected. The intima was separated and the cells were digested by trypsin.

Neurologic impairment was scored as follows. Score 0, normal activity without neurologic impairment presentation; score 1, no fully stretch in left forepaw; score 2, turning to left when walking; score 4, no spontaneous walk and loss of consciousness. The rats in score 1-3 after modeling were enrolled in this study. The rats in score 0 or 4, dyspnea, subarachnoid hemorrhage, or death after modeling were excluded.

### *The Isolation and Cultivation of Cerebrovascular Endothelial Cell*

The rat was killed and soaked in 75% ethanol. The cranial cavity was opened for the collection of cerebral cortex tissue. After washed by PBS for twice, the brain tissue was cut into pieces and digested by 0.1% collagen II at 37°C for 1.5 h. After centrifuged at 250 g for 5 min, the cell sediment was treated by 15% glucose solution. Next, the mixture was centrifuged at 4°C and 4000 rpm for 20 min. Then the cell sediment was treated by 0.1% collagen II/dispase at 37°C for 1 h and centrifuged at 1000 rpm for 5 min. At last, the cell sediment was suspended in DMEM complete medium containing 15% FBS, 1% L-glutamic acid, and 1% penicillin-streptomycin. The cells were passed at 1:4 and were used for an experiment on the 3<sup>rd</sup> generation.

### *The Establishment of VEC Oxygen-glucose Deprivation (OGD) Model*

VECs from the sham group were seeded in a six-well plate and washed by Earle's balanced salt solution twice. Then the VECs were cultured in glucose-free balanced salt solution at 95% N<sub>2</sub>, 5% CO<sub>2</sub>, and 37°C for 6 h to construct hypoxia. VECs in logarithmic phase were divided into four groups, including miR-NC, miR-124 inhibitor, pIRES2-blank, and pIRES2-PI3K groups. The cells were treated by abovementioned hypoxia at 72h after transfection.

### *Flow Cytometry*

The cells were digested by trypsin and resuspended in 500  $\mu$ L Annexin V Binding Buffer. Next, the cells were incubated in 5  $\mu$ L Annexin V-FITC and 5  $\mu$ L PI in the dark at room temperature for 15 min. At last, the cells were tested on Beckman Coulter CytoFLEX flow cytometry to evaluate cell apoptosis.

### ROS Detection

The cells were digested and incubated in 0.1% DCFH-DA probe diluted by the serum-free medium at 37°C for 20 min. After being washed for three times, the cells were tested on Beckman Coulter CytoFLEX flow cytometry.

### Dual Luciferase Activity Detection

The PCR products containing the full length of PI3K gene 3'-UTR or mutant segment were cloned to pGRE-luc. Next, it was transformed to DH5 $\alpha$  competent cells and sequenced to select the plasmid with the correct sequence. Then pGRE-PI3K-3'-UTR-wt (or pGRE-PI3K-3'-UTR-mut) was co-transfected to HEK293T cells using Lipofectamine 2000 together with miR-124 inhibitor or miR-NC. The luciferase activity was detected using the Passive Lysis Buffer and Stop&Glo solution after being cultured for 48 h.

### qRT-PCR

Total RNA was extracted using RNA Isolation Reagent and adopted for PCR reaction by QuantiTect SYBR Green RT-PCR Kit. The reaction system contained 2  $\mu$ g template RNA, 1.0  $\mu$ L primers at 0.5  $\mu$ M/L, 10  $\mu$ L 2 $\times$ QuantiTect SYBR Green RT-PCR Master Mix, 0.5  $\mu$ L QuantiTect RT Mix, and ddH<sub>2</sub>O. The PCR reaction was composed of 50°C for 5 min and 95°C for 15 min, followed by 40 cycles of 94°C for 15 s, 60°C for 30 s, and 72°C for 30 s. Real-time PCR was performed on Bio-Rad CFX96 to test the relative expression.

### Western Blot

Total protein was extracted by RIPA from cells. A total of 40  $\mu$ g protein was separated by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) and transferred to polyvinylidene difluoride (PVDF) membrane. Next, the membrane was blocked by 5% skim milk at room temperature for 60 min and incubated in primary antibody at 4°C overnight (PI3K, AKT, p-AKT, Bcl-2, and  $\beta$ -actin at 1:4000, 1:2000, 1:1000, 1:3000, and 1:10000, respectively). Then the membrane was incubated with horseradish peroxidase (HRP) labeled secondary antibody (1:30000) for 60 min after being washed by PBST for three times. At last, the protein expression was detected by enhanced chemiluminescence (ECL).

### Statistical Analysis

All data analyses were performed on SPSS 18.0 software (SPSS Inc., Chicago, IL, USA). The me-

asurement data were depicted as mean  $\pm$  standard deviation and compared by *t*-test.  $p < 0.05$  was considered as statistical significance.


## Results

### MiR-124 Inhibited PI3K Expression

The *in silico* analysis with microRNA.org online prediction showed the targeted binding site between miR-124 and 3'-UTR of PI3K mRNA (Figure 1A). Dual luciferase assay revealed that the relative luciferase activity of HEK293T cells was significantly reduced after the transfection of miR-124 mimic, while was elevated after the treatment of miR-124 inhibitor, indicating the regulatory relationship between miR-124 and PI3K mRNA (Figure 1B).


### Vascular Intima ROS Production and Cell Apoptosis were Enhanced in Vascular Intima from MCAO Rat

We, then, detected the ROS content and cell proliferation and flow cytometry assay revealed that ROS content was significantly increased in




(A)

■ miR-NC ■ miR-124 mimic  
 ■ miR-124 inhibitor



**Figure 1.** MiR-124 inhibited PI3K expression. (A) The binding site between miR-124 the 3'-UTR of PI3K mRNA; (B) Dual luciferase assay. \* $p < 0.05$ , compared with mimic NC.



**Figure 2.** Vascular intima ROS production and cell apoptosis were enhanced in vascular intima from MCAO rat. (A) Flow cytometry detection on ROS content; (B) flow cytometry detection on cell apoptosis.

vascular intima from MCAO group compared with that in sham group (Figure 2A). Moreover, cell apoptosis was aggravated in MCAO group compared with that in sham group (Figure 2B).

#### **MiR-124 level was Upregulated, while PI3K Expression was Inhibited in Vascular Intima from MCAO Rat**

The qRT-PCR demonstrated that miR-124 expression was markedly upregulated, whereas


the levels of PI3K and Bcl-2 mRNA were apparently reduced in vascular intima from MCAO rat compared with that in sham group (Figure 3A). The result of Western blot exhibited that the expressions of PI3K, p-AKT, and Bcl-2 protein were significantly alleviated in vascular intima from MCAO rat compared with that in sham group (Figure 3B).

#### **OGD Induced VECs Apoptosis, Promoted ROS Production, and Elevated miR-124 Expression**

According to the data from flow cytometry assay, we found that ROS content (Figure 4A) and cell apoptosis (Figure 4B) were markedly enhanced in VECs from OGD group compared with that in normal control. The qRT-PCR result revealed that miR-124 expression was increased, whereas the levels of PI3K and Bcl-2 mRNA were declined in VECs from OGD group compared with that in normal control (Figure 4C). Western blot further validated that the levels of PI3K, p-AKT, and Bcl-2 protein were markedly reduced in VECs from OGD group compared with that in normal control (Figure 4D).


#### **Downregulation of miR-124 or Enhancement of PI3K Attenuated the Apoptosis of VECs Induced by OGD**

After the treatment of miR-124 inhibitor or pI-RES2-PI3K transfection, the expressions of PI3K, p-AKT, and Bcl-2 were apparently enhanced (Figure 5A and B), along with the decrease of cell apoptosis (Figure 5C) and ROS content (Figure 5D) in VECs induced by OGD.



**Figure 3.** MiR-124 level was upregulated, while PI3K expression was reduced in vascular intima from MCAO rat. (A) qRT-PCR detection of gene expression; (B) Western blot detection of protein expression. \* $p < 0.05$ , compared with sham group.






**Figure 4.** OGD induced VECs apoptosis, promoted ROS production, and elevated miR-124 expression. **(A)** Flow cytometry detection on ROS content; **(B)** flow cytometry detection on cell apoptosis; **(C)** qRT-PCR detection; **(D)** Western blot detection. \* $p < 0.05$ , compared with control.

## Discussion

VECs provide a mechanical and functional barrier between blood and peripheral tissue<sup>23</sup>. Besides, VECs also exert critical functions in the regulation of blood vessel, coagulation, and fibrinolysis, through the secretion of various kinds of endothelial cell active substance<sup>3,4</sup>. VECs can produce ROS, and also could be damaged by ROS. The mediation of ROS on VECs injury thus plays a key role in the occurrence and development of ischemia stroke, hypertension, and atherosclerosis<sup>8,9</sup>. Moreover, Bcl-2 is an important anti-apoptotic factor that inhibits Cyt C release, influences calcium ionic transmembrane transport, and suppresses the activation of apoptotic protease activating factor-1 (Apaf-1)<sup>24</sup>. Of note, Bcl-2 also contributed to a crucial role in restraining ROS production<sup>25</sup>. Several studies<sup>11-13</sup> demonstrated that Bcl-2 is one of the important target genes of PI3K/AKT signaling pathway. PI3K/AKT signaling pathway plays its role in antioxidation,

antagonizing cell apoptosis, and promoting cell survival through upregulating Bcl-2 expression. This study showed that PI3K, p-AKT, and Bcl-2 levels were markedly reduced in rat VECs from MCAO group compared with sham group, which was consistent with the previous study.

Some studies<sup>20-22</sup> showed that miR-124 expression was significantly increased after ischemia stroke. Leung et al<sup>26</sup> revealed that miR-124 expression was markedly upregulated in peripheral blood from ischemia stroke patients compared with healthy control, which was correlated with the increase of NIHSS scoring. Liu et al<sup>27</sup> demonstrated that miR-124 level was abnormally increased in ischemia brain tissue from MCAO mouse. MiR-124 exhibited good diagnostic value to ischemia stroke with the areas under the curve (AUC) at 0.6976<sup>20</sup>. Consistently, we also found that miR-124 expression, ROS content, and cell apoptosis were markedly increased in rat VECs from MCAO group compared with sham group. To further investigate the potential mechanism of miR-124 on VECs, bioinformatics analysis



**Figure 5.** Downregulation of miR-124 or enhancement of PI3K attenuated VECs apoptosis induced by OGD. (A) qRT-PCR detection; (B) Western blot detection; (C) Flow cytometry detection on cell apoptosis; (D) Flow cytometry detection on ROS content. <sup>a</sup>*p* < 0.05, compared with miR-NC; <sup>b</sup>*p* < 0.05, compared with pIRES2-blank.

was performed and showed the complementary binding site between miR-124 and PI3K mRNA. Dual luciferase assay as well revealed the regulatory relationship between miR-124 and PI3K mRNA, suggesting that miR-124 might mediate the proliferation of VECs via the regulation of PI3K/AKT signaling pathway.

Sun et al<sup>21</sup> exhibited that OGD treatment markedly increased miR-124 level and apoptosis of nerve cell, suggesting that the elevation of miR-124 under hypoxia is related to cell apoptosis, which was in accordance with our results. Zhang et al<sup>28</sup> revealed that ischemia treatment apparently inhibited PI3K and AKT functions in myocar-

dium. Sayed et al<sup>29</sup> found that ischemia weakened PI3K and AKT functional activity in myocardial cells. Similar to previous data, our study also observed that the activity reduction of PI3K/AKT signaling pathway was related to ischemia hypoxia-mediated injury. Our further investigation showed that the treatment of miR-124 inhibitor or pIRES2-PI3K transfection apparently enhanced PI3K, p-AKT, and Bcl-2 expressions, along with the decrease of cell apoptosis and ROS content in VECs induced by OGD. Similarly, Ke et al<sup>30</sup> found that the upregulation of PI3K/AKT signaling pathway markedly decreased ROS production and cell apoptosis under ischemia-hypoxia

condition. Xu et al<sup>31</sup> observed that the activation of PI3K/AKT signaling pathway enhanced Bcl-2 expression and weakened cell apoptosis induced by ischemia-hypoxia. All of these studies suggested that reducing miR-124 or upregulating PI3K/AKT signaling pathway play a role in antagonizing cell apoptosis induced by hypoxia, which also supported our results. However, in our study, the regulatory mechanism of miR-124 in model rat has not been fully elucidated, which needs further investigation.

## Conclusions

MiR-124 induces the apoptosis of VECs via attenuating PI3K/AKT signaling pathway and promoting ROS production, which provides academic basis for the therapy of hypoxia in the future.

## Acknowledgments

This work was supported by Social Development Innovation Project of Xiamen Science and Technology Bureau (No. 3502Z20114022).

## Conflict of Interest

The Authors declare that they have no conflict of interest.

## References

- 1) FORSYTH MA, BARRETT T. Evaluation of polymerase chain reaction for the detection and characterisation of rinderpest and peste des petits ruminants viruses for epidemiological studies. *Virus Res* 1995; 39: 151-163.
- 2) ISMAIL TM, YAMANAKA MK, SALIKI JT, EL-KHOLY A, MEBUS C, YILMA T. Cloning and expression of the nucleoprotein of peste des petits ruminants virus in baculovirus for use in serological diagnosis. *Virology* 1995; 208: 776-778.
- 3) LIBEAU G, PREHAUD C, LANCELOT R, COLAS F, GUERRE L, BISHOP DH, DIALLO A. Development of a competitive ELISA for detecting antibodies to the peste des petits ruminants virus using a recombinant nucleoprotein. *Res Vet Sci* 1995; 58: 50-55.
- 4) MEYER G, DIALLO A. The nucleotide sequence of the fusion protein gene of the peste des petits ruminants virus: the long untranslated region in the 5'-end of the F-protein gene of morbilliviruses seems to be specific to each virus. *Virus Res* 1995; 37: 23-35.
- 5) ROMERO CH, BARRETT T, KITCHING RP, BOSTOCK C, BLACK DN. Protection of goats against peste des petits ruminants with recombinant capripoxviruses expressing the fusion and haemagglutinin protein genes of rinderpest virus. *Vaccine* 1995; 13: 36-40.
- 6) WAMWAYI HM, ROSSITER PB, KARIUKI DP, WAFULA JS, BARRETT T, ANDERSON J. Peste des petits ruminants antibodies in east Africa. *Vet Rec* 1995; 136: 199-200.
- 7) AMJAD H, QAMAR UL I, FORSYTH M, BARRETT T, ROSSITER PB. Peste des petits ruminants in goats in Pakistan. *Vet Rec* 1996; 139: 118-119.
- 8) KULKARNI DD, BHIKANE AU, SHAILA MS, VARALAKSHMI P, APTE MP, NARLADKAR BW. Peste des petits ruminants in goats in India. *Vet Rec* 1996; 138: 187-188.
- 9) NANDA YP, CHATTERJEE A, PUROHIT AK, DIALLO A, INNUI K, SHARMA RN, LIBEAU G, THEVASAGAYAM JA, BRUNING A, KITCHING RP, ANDERSON J, BARRETT T, TAYLOR WP. The isolation of peste des petits ruminants virus from northern India. *Vet Microbiol* 1996; 51: 207-216.
- 10) SHAILA MS, SHAMAKI D, FORSYTH MA, DIALLO A, GOATLEY L, KITCHING RP, BARRETT T. Geographic distribution and epidemiology of peste des petits ruminants virus. *Virus Res* 1996; 43: 149-153.
- 11) DEVIREDDY LR, RAGHAVAN R, RAMACHANDRAN S, SUBBARAO SM. Protection of rabbits against lapinized rinderpest virus with purified envelope glycoproteins of peste-des-petits-ruminants and rinderpest viruses. *Acta Virol* 1998; 42: 299-306.
- 12) SUMPTION KJ, ARADOM G, LIBEAU G, WILSMORE AJ. Detection of peste des petits ruminants virus antigen in conjunctival smears of goats by indirect immunofluorescence. *Vet Rec* 1998; 142: 421-424.
- 13) DEVIREDDY LR, RAGHAVAN R, RAMACHANDRAN S, SHAILA MS. The fusion protein of peste des petits ruminants virus is a hemolysin. *Arch Virol* 1999; 144: 1241-1247.
- 14) GUI T, SHEN K. miRNA-101: a potential target for tumor therapy. *Cancer Epidemiol* 2012; 36: 537-540.
- 15) DESJARLAIS M, DUSSAULT S, DHARI W, MATHIEU R, RIVARD A. MicroRNA-150 modulates ischemia-induced neovascularization in atherosclerotic conditions. *Arterioscler Thromb Vasc Biol* 2017; 37: 900-908.
- 16) ELIGULASHVILI R, PERL S, STRAM Y, FRIEDGUT O, SCHEICHAT N, SAMINA I, TRAININ Z. Immunohistochemical detection of peste des petits ruminants viral antigen in formalin-fixed, paraffin-embedded tissues from cases of naturally occurring infection. *J Vet Diagn Invest* 1999; 11: 286-288.
- 17) LIPKOVA J, DURIS K. The role of microRNA in ischemic and hemorrhagic stroke. *Curr Drug Deliv* 2016;
- 18) SUN AG, WANG MG, LI B, MENG FG. Down-regulation of miR-124 target protein SCP-1 inhibits neuroglioma cell migration. *Eur Rev Med Pharmacol Sci* 2017; 21: 723-729.
- 19) WU Z, HUANG W, CHEN B, BAI PD, WANG XG, XING JC. Up-regulation of miR-124 inhibits invasion and proliferation of prostate cancer cells through mediating JAK-STAT3 signaling pathway. *Eur Rev Med Pharmacol Sci* 2017; 21: 2338-2345.

- 20) HAFAR A, LIBEAU G, MOUSSA A, CECILE M, DIALLO A. The matrix protein gene sequence analysis reveals close relationship between peste des petits ruminants virus (PPRV) and dolphin morbillivirus. *Virus Res* 1999; 64: 69-75.
- 21) WAMBURA PN. Serological evidence of the absence of peste des petits ruminants in Tanzania. *Vet Rec* 2000; 146: 473-474.
- 22) DAS SC, BARON MD, BARRETT T. Recovery and characterization of a chimeric rinderpest virus with the glycoproteins of peste-des-petits-ruminants virus: homologous F and H proteins are required for virus viability. *J Virol* 2000; 74: 9039-9047.
- 23) WORRALL EE, LITAMOI JK, SECK BM, AYELET G. Xero-vac: an ultra rapid method for the dehydration and preservation of live attenuated rinderpest and peste des petits ruminants vaccines. *Vaccine* 2000; 19: 834-839.
- 24) DHINAKAR RAJ G, NACHIMUTHU K, MAHALINGA NAINAR A. A simplified objective method for quantification of peste des petits ruminants virus or neutralizing antibody. *J Virol Methods* 2000; 89: 89-95.
- 25) KONG CZ, ZHANG Z. Bcl-2 overexpression inhibits generation of intracellular reactive oxygen species and blocks adriamycin-induced apoptosis in bladder cancer cells. *Asian Pac J Cancer Prev* 2013; 14: 895-901.
- 26) LEUNG LY, CHAN CP, LEUNG YK, JIANG HL, ABRIGO JM, WANG DE F, CHUNG JS, RAINER TH, GRAHAM CA. Comparison of miR-124-3p and miR-16 for early diagnosis of hemorrhagic and ischemic stroke. *Clin Chim Acta* 2014; 433: 139-144.
- 27) LIU X, LI F, ZHAO S, LUO Y, KANG J, ZHAO H, YAN F, LI S, JI X. MicroRNA-124-mediated regulation of inhibitory member of apoptosis-stimulating protein of p53 family in experimental stroke. *Stroke* 2013; 44: 1973-1980.
- 28) ZHANG W, ZHANG JQ, MENG FM, XUE FS. Dexmedetomidine protects against lung ischemia-reperfusion injury by the PI3K/Akt/HIF-1alpha signaling pathway. *J Anesth* 2016; 30: 826-833.
- 29) ABRAHAM G, BERHAN A. The use of antigen-capture enzyme-linked immunosorbent assay (ELISA) for the diagnosis of rinderpest and peste des petits ruminants in Ethiopia. *Trop Anim Health Prod* 2001; 33: 423-430.
- 30) KE ZP, XU P, SHI Y, GAO AM. MicroRNA-93 inhibits ischemia-reperfusion induced cardiomyocyte apoptosis by targeting PTEN. *Oncotarget* 2016; 7: 28796-28805.
- 31) XU MC, SHI HM, GAO XF, WANG H. Salidroside attenuates myocardial ischemia-reperfusion injury via PI3K/Akt signaling pathway. *J Asian Nat Prod Res* 2013; 15: 244-252.