

Editorial – Obstructive sleep apnea syndrome and recurrent upper airway disease in children

D. PASSALI¹, M. RALLI²

¹Department of Medicine, Surgery and Neuroscience, ENT Clinic, University of Siena, Siena, Italy

²Department of Sense Organs, Sapienza University of Rome, Rome, Italy

Obstructive Sleep Apnea Syndrome (OSAS) and snoring are important health issues in the pediatric population, with a respective prevalence of 0.1% to 13% and 6% to 12%^{1,2} and a strict correlation with recurrent upper airway disease¹⁻³. Awareness of OSAS and snoring as health issues in children and adolescents is inadequate; there is a significant gap between the estimated number of children with OSAS and clinically diagnosed cases³. Missed or delayed treatment may put young patients at risk of growth lag, hyperactivity, attention deficit, learning disabilities, and low levels of education and literacy; there is a significant increase in the use of health services by children with OSAS⁴. Studies have demonstrated that the severity of OSAS correlates directly with total annual costs and is age-independent; other studies have shown that annual healthcare costs are reduced by one-third for children with OSAS undergoing adenotonsillectomy^{5,6}.

The clinical picture of OSAS in children includes noisy breathing, habitual snoring with or without breathing pauses during sleep, enuresis, sleeping in the sitting position, cyanosis, headache on awakening, excessive daytime sleepiness, and higher prevalence of hyperactivity attention deficit and learning disorders^{2,4,7}. Multidisciplinary approach plays a paramount role in OSAS; although a clinical suspicion can easily be formulated using a structured interview, diagnosis should be confirmed by a multidisciplinary team including pediatricians, ear-nose-throat specialists, and orthodontists², with investigation of adenotonsillar hypertrophy, craniofacial dysmorphisms, oropharyngeal abnormalities (dental malocclusions and jaw contraction), and obesity⁸, followed by polysomnography or less expensive objective testing (such as home sleep cardiorespiratory monitoring or night pulse oximetry) recently validated for diagnosis of OSAS in children⁸⁻¹⁰.

By combining the clinical profile and the results of sleep testing, and taking into account the predominant risk factors for OSAS, children can be classified into different phenotypes: a) the “classical” phenotype, a child with adenotonsillar hypertrophy with or without dental and skeletal malocclusions; b) the “adult type” phenotype, characterized by obesity and associated with aspects of the classical phenotype; c) the “congenital” phenotype, with anomalies such as micrognathia or craniofacial alterations associated with genetic syndromes. The phenotype should guide the therapeutic choice, including medical therapy (steroids and washing solutions administered as nasal spray and shower), bacteriotherapy, surgical therapy with adenoid and tonsil removal, orthodontic therapy, myofunctional treatment, and therapy with positive pressure devices².

Surgical therapy with adenotonsillectomy should be considered the first choice for children with severe OSAS and adenotonsillar hypertrophy, as indicated by clinical and objective testing criteria⁶. Immediate improvement may follow surgery, as measured by school performance and reduction of drug therapies. In the presence of comorbidities, adenotonsillectomy represents a first stage in the therapeutic program; in these cases, it is necessary to provide postsurgical follow-up in order to identify any patients who must undergo further treatment⁶.

Orthopedic-orthodontic therapy is able to reduce the symptoms and alter the natural history of OSAS. This treatment can be integrated with both medical therapy and surgical therapy. Physiotherapists, speech therapists and nutritionists contribute to the implementation of therapy and long-term management of children with OSAS.

In this supplement to the European Review For Medical and Pharmacological Sciences Journal, we invited contributions from different specialists to promote a multidisciplinary dissemination of research on OSAS, breathing-related sleep disorders, and recurrent upper airway disease in the pediatric age, with a special focus on each specialist’s approach to diagnosis and treatment of the issue and the role of recently-introduced bacteriotherapy for recurrent upper airway disease in the prevention and treatment of these conditions.

Sources of Funding

This work was supported by the Italian Society of Rhinology. The sponsor provided financial support for costs related to the publication of this article. The sponsor was not involved in the study design, in the collection and interpretation of data, in the writing of the study, or in the decision to submit the article for publication.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- 1) LUMENG JC, CHERVIN RD. Epidemiology of pediatric obstructive sleep apnea. *Proc Am Thorac Soc* 2008; 5: 242-252.
- 2) KADITIS AG, ALONSO ALVAREZ ML, BOUDEWYNNS A, ALEXOPOULOS EI, ERSU R, JOOSTEN K, LARRAMONA H, MIANO S, NARANG I, TRANG H, TSAOUSOGLOU M, VANDENBUSSCHE N, VILLA MP, VAN WAARDENBURG D, WEBER S, VERHULST S. Obstructive sleep disordered breathing in 2- to 18-year-old children: diagnosis and management. *Eur Respir J* 2016; 47: 69-94.
- 3) JOOSTEN KF, LARRAMONA H, MIANO S, VAN WAARDENBURG D, KADITIS AG, VANDENBUSSCHE N, ERSU R. How do we recognize the child with OSAS? *Pediatr Pulmonol* 2017; 52: 260-271.
- 4) MA Y, PENG L, KOU C, HUA S, YUAN H. Associations of overweight, obesity and related factors with sleep-related breathing disorders and snoring in adolescents: a cross-sectional survey. *Int J Environ Res Public Health* 2017; 4: pii: E194.
- 5) MASKI K, OWENS J. Pediatric sleep disorders. *Continuum (Minneapolis Minn)* 2018; 24: 210-227.
- 6) CHINNADURAI S, JORDAN AK, SATHE NA, FONNESBECK C, MCPHEETERS ML, FRANCIS DO. Tonsillectomy for obstructive sleep-disordered breathing: a meta-analysis. *Pediatrics* 2017; 139: e20163491.
- 7) MARCUS CL, BROOKS LJ, DRAPER KA, GOZAL D, HALBOWER AC, JONES J, SCHECHTER MS, WARD SD, SHELDON SH, SHIFFMAN RN, LEHMANN C, SPRUYT K, AMERICAN ACADEMY OF P. Diagnosis and management of childhood obstructive sleep apnea syndrome. *Pediatrics* 2012; 130: e714-755.
- 8) VILLA MP, PAOLINO MC, CASTALDO R, VANACORE N, RIZZOLI A, MIANO S, DEL POZZO M, MONTESANO M. Sleep clinical record: an aid to rapid and accurate diagnosis of paediatric sleep disordered breathing. *Eur Respir J* 2013; 41: 1355-1361.
- 9) KADITIS A, KHEIRANDISH-GOZAL L, GOZAL D. Pediatric OSAS: oximetry can provide answers when polysomnography is not available. *Sleep Med Rev* 2016; 27: 96-105.
- 10) AURORA RN, LAMM CI, ZAK RS, KRISTO DA, BISTA SR, ROWLEY JA, CASEY KR. Practice parameters for the non-respiratory indications for polysomnography and multiple sleep latency testing for children. *Sleep* 2012; 35: 1467-1473.