

Larvicidal activity of Saponin isolated from *Gymnema sylvestre* R. Br. (Asclepiadaceae) against Japanese Encephalitis vector, *Culex tritaeniorhynchus* Giles (Diptera: Culicidae)

K. ELUMALAI, S. DHANASEKARAN*, K. KRISHNAPPA*

Centre for Entamotoxicity Studies, Govt. Arts College (Autonomous), Nandanam, Chennai, Tamilnadu, India

*Unit of Pesticidal Research, Department of Zoology, Annamalai University, Annamalai Nagar, Tamilnadu, India

Abstract. – **AIM:** To determine the larvicidal activity of various extracts of *Gymnema (G.) sylvestre* against the Japanese Encephalitis vector, *Culex tritaeniorhynchus* in Tamilnadu, India. To identify the active principle present in the promising fraction obtained in Chloroform:Methanol extract of Fraction 2.

MATERIALS AND METHODS: The *G. Sylvestre* leaf extracts were tested, employing WHO procedure against fourth instar larvae of *C. tritaeniorhynchus* and the larval mortalities were recorded at various concentrations (6.25 microg/ml); the 24h LC₅₀ values of the *G. Sylvestre* leaf extracts were determined following Probit analysis. It was noteworthy, that treatment level 100 ppm exhibited highest mortality rates for the three different crude extracts and was significantly different from the mean mortalities recorded for the other concentrations.

RESULTS: The LC₅₀ values of 34.756 microg/ml (24.475-51.41), 31.351 microg/ml (20.634-47.043) and 28.577 microg/ml (25.159-32.308) were calculated in acetone, chloroform and methanol extract with the chi-square values of 10.301, 31.351 and 4.093 respectively.

CONCLUSIONS: The present investigation proved that *G. Sylvestre* could be possibly utilized as an important component in the Vector control Programme.

Key Words:

Gymnema sylvestre, *Culex tritaeniorhynchus*, Larvicidal activity, Saponin.

Introduction

Mosquitoes are vectors of many different disease agents around the world. Depending on the species, mosquitoes are vectors of protozoa (e.g. Plasmodium species) that cause malaria, the nematode worms that cause filariasis (e.g.

Wuchereria brugia) and a large number of arboviruses (e.g. yellow fever and dengue viruses) including two of great impact in the tropical and subtropical regions (i.e. yellow fever and dengue viruses). Service⁽¹⁾ recorded 2 million people primarily in tropical countries as being at risk from mosquito – borne diseases. Though, mosquito – borne diseases currently represent a greater health problem in tropical and subtropical climates, no part of the world is immune to this risk⁽²⁾. Control of mosquito – borne diseases is becoming increasingly difficult because of increasing resistance of mosquitoes to pesticides⁽³⁾. However, the discovery of the insecticide Dichloro – diphenyl – trichloroethane (DDT) in the 1940s was a major breakthrough in the control of vector borne diseases. The insecticide was highly effective for killing indoor – resting mosquitoes when it was sprayed on the walls of houses. Moreover, it was cheap to produce and active over a period of time.

The application of these chemical insecticides, which has been in use for decades, has met with setbacks in the light of the development of vector resistance and some attendant environmental hazards. Don – Pedro and Adegbite⁽⁴⁾ showed the ease with which resistance develops in their study in the health center and surroundings of the University of Lagos, where despite spraying Dichlovos insecticides on a two weekly basis for four years, a high mosquito population still persisted. In addition, Don – Pedro and Adigbite⁽⁴⁾ also reported other deleterious effects of insecticides on non – target organisms. With these problems in focus, it becomes increasingly necessary to search for an alternative in the development of environmentally safe; biodegradable, low cost, indigenous methods for mosquito control which can be used with mini-

mum care by individuals and communities in specific situations. This work aims at determining the efficacy of *Gymnema sylvestre* acetone, chloroform and methanolic leaf extracts on *Culex tritaeniorhynchus*.

Culex tritaeniorhynchus is a vector of Japanese encephalitis⁵. A human orbivirus is also been isolated from this species in China⁶. Isolates of Getah, Sindbis, Tembusu and dengue virus have also been found in this species⁷. Larvae are found in many temporary, semi-permanent and permanent ground water habitats such as rice paddies, streams, swamps, shallow marshes, low salinity tidal marshes, ponds, wells, ditches, puddles containing fresh or slightly polluted water⁷. In southern India this species is predominantly collected resting outdoors⁸. However, it has been found occurring in unusually high numbers resting indoors during the day time⁹. *C. tritaeniorhynchus* bites throughout the night with minor peaks in activity at 9 pm and 2.30 pm⁷. Females have preference cattle and pigs, but will also occasionally feed on birds and humans⁵. In india, it was observed to predominantly feed on cattle, lesser extent on humans¹⁰. Multiple feeding from two to three distinct host was observed during this study.

G. Sylvestre, rooting at nodes, leaves elliptic, acuminate, base acute to acuminate, glabrous above sparsely or densely tomentose beneath; Flowers small, in axillary and lateral umbel like cymes, pedicels long; Calyx-lobes long, ovate, obtuse, pubescent; Corolla pale yellow campanulate, valvate, corona single, with 5 fleshy scales. Scales adnate to throat of corolla tube between lobes; Anther connective produced into a membranous tip, pollinia 2, erect, carpels 2, unilocular; locules many ovuled; Follicle long, fusiform

Vector control is facing a threat due to the emergence of resistance to synthetic insecticides. Insecticides of botanical origin may serve as suitable alternative biocontrol techniques in the future. Although several plants have been reported for mosquitocidal activity, only a few botanicals have moved from the laboratory to field use, because they are poorly characterized, in most cases active principals are not determined and most of the works are restricted to preliminary screening. *G. sylvestre* is large climbers, common flora distributed in many parts of India with medicinal properties, but the larvicidal activity of this plant against the important Japanese Encephalitis vector, *C. tritaeniorhynchus* has not been reported so far. Hence, the present study was undertaken.

Materials and Methods

Extraction Procedure

The plant *G. sylvestre* was collected from the foothills of Western Ghats in Theni District, Tamilnadu, India during January –March on frequent field visit. The leaves first washed with running tap water and then shade dried on blotting paper at room temperature ($28\pm 2^{\circ}\text{C}$) for 7 days. Then the dried leaves were powdered by an electrical blender. The powder was sieved with the help of fine mesh then it was sequentially extracted in Soxhlet apparatus with acetone, chloroform and methanol. The plant material (1 kg) was loaded in the inner tube of the Soxhlet apparatus and then fitted into a round bottomed flask containing acetone. The solvent was boiled gently (40°C) by heating the mantle using the adjustable rheostat. The extraction was continued until complete extraction was achieved and the solvent in the flask was removed then substantially filled with chloroform then methanol individually. The crude extracts were then condensed with the help of rotary vacuum evaporator to yield residue (acetone 10.25 g; chloroform 8.50 g and methanol 5.25 g).

Larval Rearing

The mosquito larvae were recruited from the egg colony held at the Unit of Vector control and Phytochemical Research, Department of Zoology, Annamalai University, Annamalainagar, Tamilnadu, India, where they were reared in the laboratory. The eggs were washed with 0.01% formaldehyde solution for 30-40 minutes as recommended by Al-Masghadani et al¹¹. This is necessary as a precaution against possible microsporidian infections which might interfere with the normal development of the immature stages of mosquitoes¹² and soaked in water to facilitate hatching. After hatching, first instar larvae were distributed in bowls 30 cm in diameter and 12.5 m in depth. Care was taken to prevent overcrowding until development to early 4th instar larvae required for the study. The larvae were kept in the plastic buckets half filled with tap water and fed with dog biscuit once a day initially and twice during the later stages of development. Water in rearing container was refreshed every day by removing a little quantity of water from the rearing buckets and replacing with fresh water. This was aimed at preventing scum from forming on the water surface.

Bioassay

Larvicidal Bioassay

Standard methods for testing the susceptibility of mosquito larvae to insecticides as stipulated by WHO¹³ were followed. The bioassay were performed at a room temperature of $28 \pm 2^\circ\text{C}$, Relative humidity 75-85%, Photoperiod 12:12 (light: dark) and pH 7.0 of distilled water. Twenty five²⁵ larvae selected by means of rubber pipette were placed in 4 small separate specimen bottles containing 25 ml distilled water, and then exposed to each of the concentrations of the extracts in a final volume of 245 ml distilled water taken in 500 ml plastic bowls. This is done by tripping the contents of smaller specimen bottles into the 500 ml plastic bowls. The larvae in all the bowls were fed every twenty four hours on equal amount of dog biscuit which was spread evenly across the water surface.

Five replicates for each of the test concentration and control (without plant extracts) were tested for anti-larval effects. The larval mortalities were recorded at intervals of 12 hours. Dead larvae were identified when they failed to move after probing with a needle in the siphon or cervical region. Similar types of bioassay were conducted with different solvent extracts (concentrations of 6.25, 12.5, 25, 50, and 100 $\mu\text{g/ml}$) of the *G. sylvestre* fourth instar larval forms. Larval toxicity was also tested according to similar methodologies using the bioactive substances (from chloroform: methanol extract) isolated from thin-layer chromatographic (TLC) plates.

Phytochemical Analysis

The presence of phytochemicals was determined using the method described by Harborne¹⁴ and Evans¹⁵ with slight modifications. One or two drops of the chloroform: methanol extract were applied (using a capillary tube) to the bottom of each of the pre-coated and pre-heated (100°C for 30 min) glass plates (eight glass plates), which were prepared with silica gel G using Unoplan coating apparatus (Shadon, London, UK). After 5 minutes of drying, each of the plates was placed in the separate glass chamber for TLC analysis, with different solvent systems as the mobile phase. After the movement of solvent at the top of the plates, each plate was removed from the glass chamber and separately air-dried. After 10 minutes each of plates was sprayed with a different spraying reagent for the identification of appro-

priate phytochemical. The phytochemicals included in the study were saponins, steroid, terpenoids, flavonoids, alkaloid, essential oils, phenolics and amino acids. A qualitative test was carried out to indicate the presence of saponins¹⁶; the remaining phytochemicals were determined using TLC analysis by the application of suitable solvents and spray reagents and, in each case, R_f values were recorded.

Ultraviolet-Visual and Infrared Analysis of the Active Ingredient

The chloroform:methanol extract of the *G. sylvestre* was further chromatogrammed (25 plates) without the application of spraying reagents and each of the spots showed positive activity were separately scrapped according to their respective R_f values. Then each of the spots with their distinguishing R_f value was combined (from 25 plates) and undergoes further bioassay experiment to reveal the nature of active ingredient. As the spots exhibited positive response in Libermann Burchard reagent recorded highest larval mortality during further bioassay experiments, it undergoes spectral analysis by ultraviolet-visual (UV-Vis) and infrared (IR) spectroscopy. The UV-Vis analysis was carried out using a UV-1601 PC, SHIMADZU spectrophotometer (Shimadzu Biotech, Kyoto, Japan) with medium scan speed and sampling interval of 0.5 seconds. The IR spectroscopy analysis (Systronics India Ltd, Chennai, Tamilnadu, India) of the active spot was performed using KBr plates (JASCO FT-IR Model-420, Jasco Ltd., Tokyo, Japan) with a scanning speed of 2 mm s^{-1} . All solvents and reagents used were of analytical grade and purchased from E. Merck, India. The TLC silica gel plates (0.25 mm thickness) were prepared and equilibrated with 2% (w/w) of water before use.

Statistical Analysis

All the mortalities were counted and recorded as in percentages. The percentage mortality observed (%M) was corrected using Abbott's formula¹⁷ during the observation of the larvicidal potentiality of the plant extracts. Probit analysis¹ was used to determine the median lethal concentration LC_{50} ^{19,20}. A 3 X 5 Factorial experiment in completely randomized block design analysis of variance (ANOVA) was performed. Duncan Multiple Range Test (DMRT) was used to separate the means of the main factors (plant extract, concentration and their interactions).

Results

Larvicidal Activity

The efficacy of acetone extract of *G. sylvestre* on fourth instar larvae of *Culex tritaeniorhynchus* is presented in table.1. the larvicidal effect of the plant extract was clearly dependent on the concentrations of the extract. All the larvae maintained in the control medium survived for 24 hr, thus no mortality was observed among the control batch larvae. The experimental larvae were more susceptible to the maximum concentration hence, the maximum larval mortality of 100.00% was observed in 100 ppm concentration of acetone and methanol extracts. DMRT analysis clearly showed significant difference in the mean mortality of the larvae among the various treatment concentration of the three extracts ($p \leq 0.05$). besides, at 6.25 ppm concentration 6.88, 10.54 and 15.38% larval mortality was observed from acetone, chloroform and methanol crude extracts of *G. sylvestre*. Larvicidal activity of 20.08, 21.75, 27.92% were recorded in acetone, chloroform and methanol extracts respectively at 25.00 ppm concentration. Similar trend was also observed from the 50.00 ppm concentration of all

the three extracts. It was noteworthy that treatment level 100 ppm exhibited highest mortality rates for the three different crude extracts and was significantly different from the mean mortalities recorded for the other concentrations. The LC_{50} values of 34.756 ppm (24.475-51.41), 31.351 ppm (20.634-47.043) and 28.577 ppm (25.159-32.308) were calculated in acetone, chloroform and methanol extract with the chi-square values of 10.301, 31.351 and 4.093 respectively (Table I). In the same way the mean larval mortality obtained from various fractions were depicted in the Figure 1.

Phytochemical Analysis

The results of preliminary phytochemical analysis of the chloroform: methanol extract of the green berries of *G. sylvestre* is shown in Table II. A qualitative test indicated the presence of saponins and chromatographic analysis revealed the presence of steroids, alkaloids, terpenoids, saponins, amino acids, phenolics, flavonoids and essential oil as major phytochemicals and the absence of the terpenoids and phlorotannins following the application of different solvent systems and spraying reagents.

Table I. Lethal activities of crude extracts of *Gymnema sylvestre* at different concentrations against fourth instar larvae of *Culex tritaeniorhynchus*.

Concentrations tested ($\mu\text{g/ml}$)	Mortality (%)	LC_{50}	95% confidence limit		χ^2 value
			Lower	Upper	
Acetone extract					
Control	00.00 ^a	34.756	24.475	51.41	10.301
6.25	6.88 ^b				
12.50	20.08 ^d				
25.00	45.17 ^f				
50.00	66.83 ^h				
100.00	100.00 ⁱ				
Chloroform extract					
Control	00.00 ^a	31.351	20.634	47.043	31.351
6.25	10.54 ^b				
12.50	21.75 ^d				
25.00	52.50 ^g				
50.00	73.33 ⁱ				
100.00	99.00 ^j				
Methanol extract					
Control	00.00 ^a	28.577	25.159	32.308	4.093
6.25	15.38 ^c				
12.50	27.92 ^e				
25.00	51.17 ^g				
50.00	75.83 ⁱ				
100.00	100.00 ^j				

Values are mean \pm SD for five replications. Values not sharing a common superscript differ significantly at $p < 0.05$ (DMRT).




Figure 1. Larval mortality (%) observed in *Culex tritaeniorhynchus* 4th instar larvae tested with 3 fractions obtained from *Gymnema sylvestre* (Chloroform: Methanol in 3:7 ratio) at different concentrations.

When the isolated compounds from the TLC plates were further bio-assayed against the fourth instar larvae, the mortality was recorded in two compounds. The highest mortality (at a concentration of 100 ppm) was recorded in the first compound having $R_f = 0.818$ (94.5% in 48 hours at 100 ppm) followed by a second compound having $R_f = 0.946$ (84.33% in 48 hours at 50 ppm) with maximum absorption at 297.50 and 361.00 nm, respectively from fraction 2 during UV-Vis analysis.

Table II. Qualitative analysis of the phytochemicals of methanol extracts of *Gymnema sylvestre* tested leaves.

Phytochemicals	Compounds
Alkaloids	+
Carbohydrates	+
Saponins	+
Phytosteroids	+
Phenols	+
Phenols	+
Flavonoids	+
Terpenoids	-
Tannins	+
Phlorotannins	-

+ Presence of the phytochemical compound. - Absence of the phytochemical compound.

Discussion

Biopesticides may serve as suitable alternative to chemical insecticides in future as they are relatively safe, inexpensive and available everywhere in the world. Nowadays, mosquito control is mostly directed against larvae and only against adults when necessary. This is because the fight against adult is temporary, unsatisfactory and polluting for the environment, while larval treatment is more localized in time and space resulting in less-dangerous outcomes. Larval control can be an effective control tool due to the low mobility of larval mosquitoes, especially where the principal breeding habitats are man-made and can be easily identified²¹.

This work demonstrates the potency of *G. sylvestre* leaf extract as an effective larvicide against *C. tritaeniorhynchus* larvae; it was highly toxic to mosquito larvae. The high rates of larval mortality observed at 100 ppm within 48 hrs with LC_{50} value 28.577 ppm indicate the high toxicity of the methanol leaf extract. According to Mustafa and Al Khazraji²² *Azadirachta* showed excellent larvicidal properties at low concentrations against *Culex pipiens molestus*. Its LC_{50} value after 1 day was 62.5 µg/mL. Dua et al²³ stated that, emulsified concentration of neem oil formulation showed

95.5% reduction in larval population of *C. quinquefasciatus* in one day under field trails and thereafter 80% reduction was achieved up to the third week. The major bioactive constituents of *Gymnema sylvestre* is triterpenoid saponins (5.50%) and tannins (1.00%). Wiesman et al²⁴ reported that saponin extracted from the fruit of *Balanites aegyptiaca* showed 100% larvicidal activity against *Aedes aegypti* mosquito larvae. Results of present study are in line with earlier work done. Sharma et al²⁵ tested alcoholic and acetone extracts of *N. indium* leaf against *Anopheles stephensi* and found results of LC₅₀ at 185.99 ppm after 24 hours of exposure. Srivastava et al²⁶ examined the aqueous and methanolic extract of *N. indium* lattices against *Culex quinquefasciatus* and obtained that different dilutions of the lattices delay the post embryonic development of *Culex larvae*, methanolic extract is 1.8 times more toxic than aqueous extract. Datura metel showed less activity among the tested plant extracts with 54% mortality at 5 ppm.

The secondary compounds of plants make up a vast repository of compounds with a wide range of biological activities²⁷⁻³¹. Most studies report active compounds as steroidal saponins. Saponins are freely soluble in both organic solvents and water, and they work by interacting with the cuticle membrane of the larvae, ultimately disarranging the membrane, which is the most probable reason for larval death³². Wiesman and Chapagain²⁴ reported that saponin extracted from the fruit of *Balanites aegyptiaca* showed 100% mortality against larvae of *S. aegypti*. The larvicidal property of a saponin mixture isolated from *Cestrum diurnum* was also evaluated against *Anopheles stephensi* mosquito by Ghosh and Chandra³³. Recently, Krishnappa and Elumalai³⁴ reported that essential oil obtained from *Clauseana excavata* exhibited strong larvicidal and ovicidal activity against *Spodoptera litura*. These findings have re-emphasised the need to explore the possibility of using plant based larvicide and reduce the chemical hazards in the environment. The leaf extract of *G. sylvestre* was very promising. Furthermore, all these plant materials can be easily collected from the nature. Therefore, plant originated insecticides can be used as sustainable larvicide in the mosquito control programme.

The present investigation revealed that the presence of several bioactive compounds in *G. sylvestre*. However, the IR spectra of the bioactive compounds during the present study also indicated that any saponin compound(s) is responsible for larval toxicity.

Conclusions

G. sylvestre offers promise as potential bio control agent against *C. tritaeniorhynchus* particularly in its markedly larvicidal effect. The extract or isolated bioactive phytochemical could be used in stagnant water bodies for the control of mosquitoes acting as vector for many communicable diseases. The extract or isolated bioactive phytochemical from the plant could be used in stagnant water bodies which are known to be the breeding grounds for mosquitoes. However, further studies on the identification of the active principals involved and their mode of action and field trials are needed to recommend *G. sylvestre* as an anti-mosquito product used to combat and protect from mosquitoes in a Vector Control Program.

Acknowledgements

We are grateful to Professor N. Kalaichelvi, Zoology, Department of Advanced Zoology and Biotechnology, Govt. Arts College (Autonomous), Nandanam, Chennai-35 for laboratory facilities provided. Professor Dr. Govindaraju, Department of Chemistry, Kongu Polytechnique College, Erode, Tamilnadu, India, for his contribution in analysis and interpretation of data. We extend our gratitude to Dr Anbalagan, Taxonomist, Department of Botany, Annamalai University, Annamalaiagar, Tamilnadu, India for the taxonomic identification of the selected plant.

Conflict of Interest

None to declare.

References

- 1) SERVICE MW. Handbook on the African-tropical Toxorhynchitine and *Culex* mosquitoes excerpting *Aedes* and *Culex*. Brit Mus Nat Hist, London 1990; p. 207.
- 2) FRADIN MS, DAY JF. Comparative efficacy of insect repellents against mosquito bite. N England J Med 2002; 347: 13-18.
- 3) RANSON H, ROSSITER L, ORTELLI F, JENSEN B, WANG Z, ROTH CW, COLLINS FH, HEMINGWAY J. Identification of a novel class of insect glutathione S-transferases involved in resistance to DDT in the malaria vector *Anopheles gambiae*. Biochem J 2001; 359: 295-304.
- 4) DON PEDRO KN. The effects of insecticides on non-target arthropods in cowpea ecosystem. M.Sc.Thesis submitted to University of Ibadan, Nigeria 1976.

- 5) BRAM RA. Contributions to the mosquito fauna of Southeast Asia (Diptera: Culicidae) II. The genus *Culex* in Thailand. Cont Amer Entomol Inst 1967; 2: 1-296.
- 6) ATTOUI H, JAAFAR FM, BELHOUCHE M, ALDROVANDI N, TAO S, CHEN B, LIAN G, TESH RB, DE MICCO P, DE LAMBALLERIE X. Yuman orbivirus, a new orbivirus isolated from *Culex tritaeniorhynchus* mosquitoes in China. J Gen Virol 2005; 86: 3409-3417.
- 7) LEE DJ, HICKS MM, DEBENHAM ML, GRIFFITHS M, MARKS EN, BRYAN JH, RUSSELL RC. The culicidae of the Australian region. 1989; Vol. 7. Canberra, Australian Government publishing Service.
- 8) DAS BP, LAL S, SAXENA VK. Outdoor resting preference of *Culex tritaeniorhynchus*, the vector of Japanese encephalitis in Warangal and Karim Nagar districts, Andhra Pradesh. J Vect Borne Dis 2004; 41: 32-36.
- 9) KANOJA PC, GEEVARGHESE G. First report on high-degree endophilism in *Culex tritaeniorhynchus* (Diptera:Culicidae) in an area endemic for Japanese encephalitis. J Med Entomol 2004; 41: 994-996.
- 10) ARUNACHALAM N, PHILIP SAMUEL P, HIRIYAN J, RAJENDRAN R, DASH AP. Observation of the multiple feeding behaviour of *Culex tritaeniorhynchus* (Diptera:Culicidae), the vector of Japanese encephalitis in Kerala in Southern India. Am J Trop Med Hyg 2005; 72: 198-200.
- 11) AL-MASHHADANI HM, DAVIDSON G, CURTIS C. A genetic study of the susceptibility of *Anopheles gambiae* to *Plasmodium bergeri*. Trans Roy Soc Tropical Med Hyg 1980; 749: 585-594.
- 12) ANOSIKE JC, ONWULIRI COE. Experiment on *Wuchereria bancrofti* infection of *Culex quinquefasciatus* and *Aedes aegypti*. Angew Parasitol 1992; 33: 139-142.
- 13) WHO. Report of the WHO Informal Consultation on the evaluation and testing insecticides. CTD/WHOPES/IC 1996; p. 69.
- 14) HARBORNE JB. Phytochemical Methods, A Guide to Modern Techniques of Plant Analysis. London: Chapman and Hall 1984; pp. 49-188.
- 15) EVANS WC. Trease and Evans Pharmacognosy (15th edition), W.B Saunders Company Ltd 2002; pp. 135-150.
- 16) KOKATE A. Phytochemical methods. Phytotherapy 1999; 78: 126-129.
- 17) ABBOTT WS. A method for computing the effectiveness of an insecticide. J Econ Entomol 1925; 18: 265-267.
- 18) FINNEY DJ. Probit analysis (Third edition). 3rd edition. Cambridge University Press, UK 1971; p. 38.
- 19) WOOLF CM. Principles of Biometry: Statistics for Biologists. D. Van Nostrand Company Inc 1968; pp. 1-267.
- 20) WARDLAW HC. Practical Statistic for Experimental Biology. John Wiley and Sons, New York; 2002; pp. 256-268.
- 21) HOWARD AFB, ZHOU G, OMLIN FX. Malaria mosquito control using edible fish in western Kenya: preliminary findings of a controlled study. BMC Public Health 2007; 7: 199-204.
- 22) MUSTAFA MA, AL KHAZARAJI A. Effect of some plant extracts on the *Culex pipiens molestus* Forskal larvae. Iraqi J Vet Sci 2008; 22: 9-12.
- 23) DUA VK, PANDEY AC, RAGHAVENDRA K, GUPTA A, SHARMA T, DASH AP. Larvicidal activity of neem oil (*Azadirachta indica*) formulation against mosquitoes. Malaria J 2009; 8: 120-124.
- 24) WIESMAN Z, CHAPAGAIN BP. Larvicidal effects of aqueous extracts of *Balanites aegyptiaca* (desert date) against the larvae of *Culex pipiens* mosquitoes. Afr J Biotechnol 2005; 4: 1351-1354.
- 25) SHARMA P, MOHAN L, SRIVASTAVA CN. Larvicidal potential of *Nerium indicum* and *Thuja orientalis* extracts against malaria and Japanese encephalitis vector. J Environ Biol 2005; 26: 657-660.
- 26) SRIVASTAVA VK, SINGH SK, RAI M, SINGH A. Toxicity of *Nerium indicum* and *Euphorbia royleana* latices against *Culex quinquefasciatus* mosquito larvae. Nigerian J Nat Prod Med 2003; 7: 61-64.
- 27) ELUMALAI K, DHANASEKARAN S, VEERAYAN G, MEENAKSHI V. FEEDING deterrent activity of certain plant extracts against the fruitborer, *Helicoverpa armigera* (Hubner). Appl Zool Res 2008; 19: 133-138.
- 28) ANANDAN A, KRISHNAPPA K, MATHIVANAN T, ELUMALAI K, GOVINDARAJAN M. Bioefficacy of *Hyptis suaveolens* and *Melochia chorcorifolia* against the armyworm, *Spodoptera litura* (Fab.) (Lepidoptera: Noctuidae). Int J Curr Res 2010; 4: 117-121.
- 29) ELUMALAI K, KRISHNAPPA K, ANANDAN A, GOVINDARAJAN M, MATHIVANAN T. Antifeedant activity of medicinal plant essential oils against *Spodoptera litura* (Lepidoptera:Noctuidae). Int J Rec Sci Res 2010; 2: 62-68.
- 30) ELUMALAI K, KRISHNAPPA K, ANANDAN A, GOVINDARAJAN M, MATHIVANAN T. Larvicidal and ovicidal activity of seven essential oil against lepidopteran pest *Spodoptera litura* (Lepidoptera:Noctuidae). Int J Recent Sci Res 2010; 1: 8-14.
- 31) ANANDAN A, KRISHNAPPA K, GOVINDARAJAN M, ELUMALAI K. Antifeedant activity of some plant extracts against the fourth instar larvae of *Spodoptera litura* (fab.). Int J Rec Sci Res 2011; 2: 1-3.
- 32) HOSTETTMANN K, MARSTON A. Saponins (Chemistry and Pharmacology of Natural Products). Cambridge: Cambridge University Press; 1995: pp. 1-132.
- 33) GHOSH A, CHANDRA G. Biocontrol efficacy of *Cestrum diurnum* (L.) (Solanales: Solanaceae) against the larval forms of *Anopheles stephensi*. Nat Prod Res 2006; 20: 371-379.
- 34) KRISHNAPPA K, ELUMALAI K. Chemical composition of essential oil from *Clausena excavata* and its larvicidal and ovicidal activities against armyworm, *Spodoptera litura* (Fab.) (Lepidoptera: Noctuidae). Asian J Trop Dis 2012; 1-6.