

# Author Correction: Waist circumference and the risk of lumbar and femur fractures: a nationwide population-based cohort study

G.R. PARK, H.S. KIM, Y.T. KIM, H.J. CHUNG, S.J. HA, D.W. KIM, D.R. KANG, J.Y. KIM, M.Y. LEE, J.Y. LEE

Department of Internal Medicine, Yonsei Wonju College of Medicine, Wonju, Republic of Korea

Correction to: European Review for Medical and Pharmacological Sciences 2021; 25 (3): 1198-1205–DOI: 10.26355/eurrev\_202102\_24822–PMID: 33629289, published online 15 February, 2021.

After publication, the authors requested to correct Table I and some minor errors in the text.

The Publisher apologizes for any inconvenience this may cause.

The correct sections are reproduced below.

- In the abstract section, the logistic regression analysis has been changed to Cox regression analysis.
- In the results section, tertial has been replaced with tertile.
- In Table I, the term “women” has been moved to the right column.
- The caption has been added to Table I. “All values are presented as number (%) or means  $\pm$  SDs. BMI = Body mass index; SBP = Systolic blood pressure; DBP = Diastolic blood pressure; HDL = High density lipoprotein; ALT = Alanine aminotransferase; AST = Aspartate aminotransferase; GGT = Gamma-glutamyltransferase; GFR = Glomerular filtration rate. aHeavy drinker was defined as consuming seven or more drinks on the same occasion, more than 3 days per week. bHeavy exercise was defined as moderate or heavier regular physical exercise performed at least 30 minutes a day, on 3 or more days per week”.
- Part of the caption of Table II has been changed “Model 2 was adjusted for the variables in model 1. Model 3 was adjusted for the variables in model 2...”.
- Part of the caption of Table III has been changed “Model 2 was adjusted for the variables in model 1. Model 3 was adjusted for the variables in model 2...”.