

Performance improvement in sport through vitamin D – a narrative review

K. WEISS¹, A. DEVRIM-LANPIR², Z. JASTRZĘBSKI³, P.T. NIKOLAIDIS⁴, J. HILL⁵,
B. KNECHTLE^{1,6}

¹Institute of Primary Care, University of Zurich, Zurich, Switzerland

²Department of Nutrition and Dietetics, Faculty of Health Sciences, Istanbul Medipol University, Istanbul, Turkey

³Gdansk University of Physical Education and Sport, Gdańsk, Poland

⁴School of Health and Caring Sciences, University of West Attica, Athens, Greece

⁵Department of Pediatrics, Division of Gastroenterology and Nutrition, McMaster University, Hamilton, Canada

⁶Medbase St. Gallen Am Vadianplatz, St. Gallen, Switzerland

Abstract. – OBJECTIVE: It is well known that vitamin D deficiency can lead to various health problems. However, it is not common knowledge among athletes and doctors that vitamin D deficiency is prevalent in sports. This deficiency can severely impact performance, while vitamin D supplementation can alleviate this effect and potentially improve performance.

MATERIALS AND METHODS: This narrative review aims to compile the current state of knowledge about the importance of vitamin D in increasing performance for active people. To this end, we searched the 'Scopus' and 'PubMed' databases for the terms 'vitamin D athlete – performance' with an end date of 31/12/2022.

RESULTS: Study results indicated the therapeutic impact of vitamin D on aerobic capacity, recovery, strength and sprint performance remains controversial.

CONCLUSIONS: Based on previous findings, 4,000-5,000 IU of vitamin D per day may be a safe dose that can improve athletic performance.

Key Words:

Vitamin D Deficiency, Aerobic capacity, Strength performance, Recovery.

Introduction

It is well known that a vitamin D deficiency can lead to various health problems. Vitamin D refers to the compound cholecalciferol and, when synthetically derived, calcitriol¹. Historically, vitamin D is an essential component of calcium and phosphate homeostasis and metabolism within bones and skeletal muscles². It has also been implicated in

several other physiological functions throughout the body³. There are two primary forms of vitamin D in humans, cholecalciferol (vitamin D₃, from animal sources) and ergocalciferol (vitamin D₂, from plant sources), both of which can be ingested through food, supplementation, or sun exposure⁴.

Risk factors of a vitamin D deficiency have been well described in the general population⁵⁻⁷. Numerous studies⁸⁻¹² on vitamin D levels in various groups of athletes, including runners, basketball players, jockeys, gymnasts and even dancers, have been conducted in the last decade, showing that they were comparable to those of the general population. However, athletes and doctors are unlikely to be aware that vitamin D deficiency is prevalent in sports^{13,14} depending on the season^{15,16}. This deficiency can negatively affect muscle strength, power, and endurance¹³. Furthermore, low vitamin D levels can increase the occurrence of stress fractures and other musculoskeletal injuries and can affect acute muscle injuries and inflammation following high intensity exercises¹³. Vitamin D deficiency can severely limit athletic performance¹⁷⁻¹⁹, but it has been shown that vitamin D supplementation can alleviate this effect and improve performance²⁰⁻²².

Multiple reviews^{13,16,23,24}, about the potential ergogenic effects of vitamin D on athletic performance have been published. In 2015, the aspects of skeletal muscle function, increased force and power production combined with decreased recovery times were studied²³. In 2020 the aspects of the immune system and musculoskeletal function were considered¹⁶. Again in 2020, the aspects of muscular strength, cardiovascular health and the incidence of illness, including upper respira-

tory tract infections, were investigated²⁴, and in 2021 physical performance and musculoskeletal injuries in athletes were considered¹³. At the time of writing, there is no review exploring vitamin D deficiency in endurance and strength athletes and the potential ergogenic effects of vitamin D supplementation. Therefore, this narrative review aims to compile the current research about the importance of vitamin D and its impact on the performance of endurance and strength athletes.

Materials and Methods

This review utilized a narrative review method described by Grant and Booth²⁵. We searched two significant databases (Scopus and PubMed) using terms related to vitamin D and athletic performance (vitamin D; athletic performance; sports performance; athlete; professional athlete; elite athletes; college athletes; sports; performance improvement) for articles published before July 2022. The authors sorted the search results (case reports, original papers and review articles) and were included in this review based on their clinical relevance to athletes and practitioners.

Definition of Vitamin D Deficiency

An essential factor to consider when determining vitamin D levels is the classification of a limit for a deficiency (Table D). There has been a significant debate around the definition of deficiency and insufficiency in recent years. Vitamin D deficiency is generally divided into vitamin D insufficiency (50-75 nmol/l), vitamin D deficiency (<50 nmol/l) and severe vitamin D deficit (<25 nmol/l)²⁶. Elsewhere, a vitamin D level of <50 nmol/l and a vitamin D insufficiency of 50-75 nmol/l are reported²⁷, although the recommendations highlight the importance of vitamin D, consequently statements differ depending on the specific society or regulatory body¹. Therefore, it is essential to consider particular countries' guidelines when determining vitamin D deficiency.

Another significant issue arising from various recommendations is the measurement units used. The classification uses nmol/l and others ng/ml. In the past, vitamin D levels of <10 ng/ml were defined as a deficit because the serum level of vitamin D and calcium absorption decreased significantly below this threshold^{28,29}. However, the IOM³⁰ defines vitamin D insufficiency as a serum concentration of 25 hydroxyvitamin D [25(OH)D] <30 ng/ml (<75 nmol/l). Elsewhere, the vita-

min D deficit is defined with a serum concentration of 25(OH)D <20 ng/ml (<50 nmol/l) and vitamin D insufficiency with a serum concentration of 25(OH)D <30 ng/ml (<75 nmol/l)³¹. Since the classification varies, we have used the term vitamin D deficiency unless otherwise described.

There is inconsistency in the classification of a deficiency. Although vitamin D deficiency is generally divided into two categories, vitamin D insufficiency and vitamin D deficiency, different scientific societies and national agencies refer to the cut-off values of serum vitamin D concentrations differently (Figure 1). For example, the National Academy of Medicine and the American College of Medicine reported serum 25(OH)D concentration below 12 ng/ml (<30 nmol/L) as vitamin D deficiency and between 10-30 ng/mL (<30 nmol/L) as vitamin D insufficiency. At the same time, the European Society of Endocrinology defines serum 25(OH)D concentration below <20 ng/ml (<50 nmol/L) as vitamin D deficiency and between 20-30 ng/ml (<50-75 nmol/L) as vitamin D insufficiency³². For this reason, different cut-off values were used when interpreting the results of studies on vitamin D and athletic performance^{9,17,18}. In addition, a comprehensive review of vitamin D in athletes recommended 30 ng/mL (75 nmol/L) as the cut-off value for vitamin D adequacy when assessing serum concentrations in athletes before vitamin D supplementation³³.

Vitamin D and Athletic Performance

In recent years vitamin D was suggested to have ergogenic properties that may improve sports performance. Willis et al³⁴ recently highlighted that a surprisingly high percentage of athletes are probably vitamin D deficient, especially those exercising an indoor sport. New studies³⁵⁻³⁸ clearly show a connection between vitamin D status and performance. However, this relationship could be gender-specific and age-dependent³⁸ favoring young men¹⁷. A recent double-blind study³⁹ in non-diabetic men found that supplementation of 3,332 IU/day of vitamin D significantly increased the serum levels of 25(OH)D, total testosterone, bioactive testosterone, and free testosterone levels.

Vitamin D₃ receptors (VDR) are found in human skeletal muscle tissue^{40,41}, suggesting that 1,25-dihydroxy vitamin D may directly influence skeletal muscle activity. Furthermore, detecting the VDR in the muscles indicated that vitamin D may have an essential function in them^{42,43}. The question arose as to whether a lack of vitamin D could influence skeletal muscle damage caused by overuse⁴³.

Studies on the effects of vitamin D₃ in muscles²¹ have been limited primarily to population groups with diseases⁴⁴ or healthy, untrained adults⁴⁵. Until recently, studies⁴⁴⁻⁴⁶ have shown that increasing serum 25(OH)D levels in specific populations positively affects muscle strength and muscle mass. Contrary to those findings, the only study that investigated these effects in athletes showed different results⁴⁷. In addition, von Hurst and Beck⁴⁷ concluded that the optimal dosage of 25(OH)D serum concentrations in athletes cannot be conclusively assessed yet.

The Importance of Vitamin D on Maximum Oxygen Uptake

In athletes, the effect of vitamin D was mainly investigated concerning maximum oxygen uptake (VO₂max)⁴⁸⁻⁵¹, endurance⁵² and strength performance⁵³⁻⁵⁶. While VO₂max, endurance and strength performance seemed to be positively influenced, the anaerobic performance was not affected⁵⁷.

The mechanism that acts on VO₂max when serum 25(OH)D levels are increasing is still unclear⁵⁰. However, this could be explained by the CYP enzymes (cytochrome P), which activate vitamin D₃ to 1,25-dihydroxy vitamin D₃ and bind it to heme-containing proteins⁵⁸ and could influence the binding affinity of oxygen to hemoglobin. In addition, VDRs are present in the myocardium and vascular tissue⁵⁹, suggesting that 1,25-dihydroxy vitamin D may influence VO₂max via the ability to transport and increase oxygen in the blood. Several studies^{48,50} have shown a positive association between VO₂max and serum 25(OH)D concentration in athletes. However, additional variables were not considered, such as the simultaneous use of multivitamin preparations⁴⁸ and dietary supplements^{50,60}.

Studies on the relationship between aerobic power (determined by VO₂max) and vitamin D in athletes are contradictory. Tsouros et al⁶¹ found a significant correlation between 25(OH)D values and selected performance parameters in male professional footballers. A linear relationship was seen between 25(OH)D values before and after the season with measurements of isometric strength and VO₂max⁶¹. In another study⁶², young soccer players received either vitamin D (5,000 IU/day) or a placebo during eight weeks of intensive training (high-intensity interval training, HIIT). In the supplementation group, vitamin D concentration increased by 10% and VO₂max by 20%; in the placebo group, vitamin D concentration decreased by 8.4% and

VO₂max increased by 13%, suggesting that vitamin D supplementation may have a moderate, though not significant, impact on aerobic capacity in youth soccer players.

In another study⁶³, lightweight rowers with sufficiently high vitamin D levels (>30 ng/ml) were observed during an eight-week training phase where either 6,000 units of vitamin D or placebo were administered daily. It came to a significant increase in VO₂max (12.6% compared to 10.3%) and an increase in vitamin D levels around 100%⁶³. On the other hand, a more recent study showed no correlation between 25(OH)D and VO₂max in ice hockey players. In addition, the relationship usually observed between the serum concentration of 25(OH)D and VO₂max is inversely correlated with increased physical activity and training regimen⁵⁰. Similar interactions were observed in Polish elite footballers³⁰ and young Russian footballers¹⁹. In addition, a study of Gaelic football players⁵¹ stated that even 50% of participants had inadequate levels of vitamin D. The daily intake of 3,000 IU vitamin D₃ for 12 weeks only corrected the vitamin D deficiency but had no meaningful effect on VO₂max⁵¹. Further studies are needed to explore this interaction and verify whether athletes with a deficiency of 25(OH)D or an ergogenic effect exist and whether it is the supraphysiological doses of vitamin D₃, as used by Jastrzębski⁶³, that have an ergogenic effect.

The Importance of Vitamin D on Strength Performance

Recent studies show several connections between vitamin D levels and strength performance in healthy individuals^{37,45,53-56,64-66} and athletes^{37,45,53-56,64-66}. It is suggested that vitamin D₃ may increase the strength and performance of skeletal muscle tissue^{43,67}, possibly through sensitization of the calcium-binding sites on the sarcoplasmic reticulum, which leads to increased muscle contractions⁶⁸. There is also further evidence that vitamin D₃ may increase the size and the number of type II muscle fibres⁶⁹⁻⁷¹ and improve muscle function through morphological adjustments and improved calcium availability⁶⁹. However, these findings have yet to be verified in studies with athletes.

So far, it seems questionable whether low vitamin D levels correlate to poor performance and whether consuming high doses of vitamin D improves performance. Cross-sectional studies indicated that vitamin D concentrations were associated with muscular performance in col-

Table 1. Summary of the outcomes from controlled studies investigating the potential efficacy of Vit D supplementation on performance in athletes.

Country	Study design	N (men/ women)/age Participants	Participants	Baseline 25(OH)D	Vitamin D dose/duration	25(OH)D concentrations after intervention	Outcomes	Ref.
USA	RCT	19 (6/13) / 20.1±1.6 yrs (Vit D) - 19.7±0.8 yrs (PLA)	College Swimmers	58±21.75 nmol/L	5,000 IU Vit D3/12 wks	8% increase in Vit D group; 44% decrease in PLA group	↑ in squat, vertical jump, and deadlifts ↔ in bench press, standing long jump, dips and pull-ups performance	114
Tunisia	RCT	36 (36/0) / 10.7±2.15 yrs (Vit D) - 10.8±12.2 yrs (PLA)	Recreational footballers	12.4±3.49 ng/ml (Vit D) - 12.1±3.93 (PLA)	200,000 IU Vit D3/ a bolus dose	28.6±5.34 ng/ml (Vit D) - 19.3±4.1 (PLA)	↑ in vertical jump, triple hop, 10m/30m sprints and shuttle run, ↔ in long jump while standing	115
UK	RCT	30 (30/0) / 22±2 yrs (20,000 IU/wk group); 21±1 yrs (40,000 IU/wk group); 21±1 yrs (PLA)	Club-level athletes (football, rugby)	53±26 nmol/L (20,000 IU/ wk group); 51±26 (40,000 IU/ wk group); 52±27 nmol/L (PLA)	20,000 or 40,000 IU Vit D3 per wk/12 wks	Supplementation with 40,000 IU Vit D induced greater values than 20,000 IU Vit D	↔ in physical performance parameters	80
UK	Controlled prospective study	24 (11/5) / 26±4.57 yrs	Classic Ballet dancers	NS	2,000 IU Vit D3/ 4wks	NS	↑ in isometric strength and vertical jump	116
Brazil	Single blind/ Crossover study	27 (27/0) / 24±4 yrs	Combat sports athletes (MMA, Jiu-jitsu and boxers)	-	50,000 IU/ 80,000 IU/ 100,000 IU/ 150,000 IU/ 200,000 IU/ 250,000 IU/ 300,000 IU/ 350,000 IU/ 400,000 IU/ 450,000 IU/ 500,000 IU/ 6 weeks per group	-	↑ in upper and lower body VO ₂ peak, upper body Wingate performance - Suppl with more than 50,000 IU vit D showed no additional benefit	117
Spain	RCT	36 (36/0) / 27±6 yrs	Elite rowers	26.24±8.18 ng/ml (Vit D) - 26.95 ng/ml (PLA)	200,000 IU Vit D3/ 6 weeks	48.12±10.88 ng/ml (Vit D); 35.14±7.96 ng/ml	↔ in muscle recovery	96
UK	RCT	10 (10/0) / average age of 18 yrs	Soccer players	Presented in figure/ Numerical data NS	5,000 IU Vit D3/ 8 weeks	Serum vit D cons increased in Vit D group	↑ in 1-RM bench press and 1-RM back squat, ↔ in 30-m sprint and Illinois agility run	54
Iran	RCT	69 (36/33) / 18-23 yrs	Handball and futsal athletes	27.5±17.9 ng/ml (Vit D); 27.5±17.9 ng/ml (PLA)	50,000 IU Vit D3 per week/ 8 weeks	44.9±19.8 ng/mL (Vit D); 21.3±8.9 ng/mL (PLA)	↑ in the strength leg press test and sprint test, ↔ in the ergo jump test, vertical jump test, Illinois agility test, 40-yard speed test	72

Continued

Table I. Summary of the outcomes from controlled studies investigating the potential efficacy of Vit D supplementation on performance in athletes.

Country	Study design	N (men/ women)/ age	Participants	Baseline 25(OH)D	Vitamin D dose/duration	25(OH)D concentrations after intervention	Outcomes	Ref.
UK	RCT	22 (22/0)/29±10 yrs (Vit D)- 26±7 (placebo)	Elite indoor athletes (Judokas)	13.16±3.75 ng/mL (Vit D); 16.33±2.73 ng/mL (PLA)	150,000 IU a bolus dose	16.76±3.21 ng/mL (Vit D); 16.33±2.56 ng/mL (PLA)	↑ in muscle strength seven days after supplementation	118
New Zealand	Prospective, double blind, placebo-controlled study	57 (57/0)/ average age of 21.5 yrs (Vit D)- 20.9 yrs (placebo)	Rugby players	93±19nmol/L (Vit D); 95±17 nmol/L (PLA)	50,000 IU biweekly/ 12 weeks	111±18 nmol/L (Vit D-wks 5-6); 114±19 nmol/L (Vit D-wks 11-12); 85±17 (PLA-wks 5-6); 81±21 (PLA-wks 11-12)	Improvement of six physical performance parameters including 30m sprint time; ↑ in the weighted reverse-grip chin up performance	82
South Korea	RCT	35(21/14)/ 20.1±0.15 yrs	Collegiate Taekwondo athletes	27.3±1.18 nmol/L (Vit D); 30.9±1.95 nmol/l (PLA)	5,000 IU/ 4 weeks	91.7±2.4 nmol/l (Vit D); 87.7±2.4 nmol/l (PLA)	↑ in peak performance and isokinetic knee extension at 180° ↔ in Wingate test, endurance, counter-movement jump, sit-ups, agility test, 20m pacer)	119
Poland	RCT	36 (36/0)/ average age of 17-19 yrs	Football players	48.5±8.6 nmol/L (Vit D); 47.5±16.2 nmol/L (placebo)	5,000 IU Vit D3/8 weeks	71.5±12.7 nmol/l (Vit D); 57.8±21.7 nmol/l (PLA)	↑ in serum Vit D; ↔ in Wingate test, 5-, 10-, 20-, 30-m sprints, knee bend jumps, counter-movement jumps, small field games, interval running at AT	81
Ireland	Double-blind, placebo-controlled study	42 (18/24)/ average age of 20±2 yrs	Gaelic football players	47.37±13.29 nmol/L (Vit D); 43.10±10.00 (PLA)	3,000 IU Vit D/ 12 weeks	83.68±32.98 nmol/l (Vit D); 49.22±25.40 nmol/l (PLA)	No change in physical performance (VO ₂ max, vertical jump height, left and right handgrip strength, forced vital capacity, forced expiratory volume at 1s)	51
Poland	RCT	36 (sex was not defined) average age of 17.5±0.6 yrs	Football players	45±8.5 nmol/l (Vit D); 47.5±16.2 nmol/l (PLA)	6,000 IU Vit D/ 8 weeks	106.3±26.6 mmol/l (Vit D); 42.5±16.7 mmol/l (PLA)	↑ aerobic capacity in both groups (20% in the Vit D group; 13% in the PLA group)	62
Poland	RCT	28 (sex and age were not defined)	Elite football players	32.37±6.7 nmol/l (Vit D); 25-30 ng/ml (PLA)	6,000 IU/ 6 weeks	54.03±7.4 ng/ml (Vit D); 35-40 ng/ml (PLA)	Improvement in 5-m sprint, no change in VO ₂ max and 30-m sprint performance	120




Figure 1. Different scientific societies develop diverse sets of 25(OH)D concentrations [ng/mL]. NAM: National Academy of Medicine, US; ES: Endocrine Society, US; ESE: European Society for Endocrinology, UK; EFSA: European Food Safety Authority, Europe; SACN: Scientific Advisory Committee on Nutrition, UK.

legiate athletes³⁷, UK-based athletes³⁸, young Danish elite swimmers^{37,45,53-56,64-66}, and skeletal muscle strength and muscle power in young Polish elite judokas^{37,45,53-56,64-66}. In addition, a randomized, controlled trial⁷² of Romanian athletes who received 50,000 IU of vitamin D per week for eight weeks indicated a meaningful improvement in strength performance (leg press). However, inconsistent with those studies, no association was observed between vitamin D serum concentrations and skeletal muscle performance in studies with Polish elite footballers⁴⁹, young Russian footballers⁵⁰, Swedish female soccer players⁷³, and Gaelic footballers⁵¹.

Data on muscle strength and athletic performance in athletes is limited, and results are controversial. The few studies^{14,74} that explored the interaction between muscle strength and serum vitamin

D status in athletes found no significant association between them. A comparison between athletes with sufficient (>50 nmol/l) and athletes with insufficient (<50 nmol/l) vitamin D status showed no differences in counter-movement jump and aerobic fitness (VO₂max)⁷⁴. Similar findings were reported in a study investigating male professional basketballers. In athletes with deficient (<20 ng/ml), insufficient (20-32 ng/ml), and sufficient vitamin D (>32 ng/ml), muscle strength of knee extension/flexion and ankle dorsiflexion/plantarflexion was investigated. Interestingly, the isokinetic lower limb strength was not significantly different when all three groups were compared¹⁴.

It is stated that although vitamin D may have beneficial effects on skeletal muscles in athletes, it has no effect on the performance of athletes that do not exhibit a vitamin D deficiency⁷⁵. However,

some studies^{64,76,77} on muscle strength and vitamin D in athletes showed no effect on performance even though the vitamin D level was deficient. For example, in the study⁶⁴ examining hockey players with vitamin D deficiency, no relationship was found between serum vitamin D levels and physiological performance variables. Another study⁷⁶ investigated an association between vitamin D and lower extremity muscle strength in 342 football players. Although 84% of the athletes had vitamin D deficiency, no correlation was detected between low vitamin D levels and lower extremity muscle strength. A study⁷⁷ including 80 competitive swimmers (42 vitamin D insufficient and 11 vitamin D deficient) found no association between vitamin D levels and various performance parameters. In particular, there were no differences between athletes with and without vitamin D deficiency.

In some studies, muscle function⁴⁷ and athletic performance⁷¹ were only improved in athletes with a vitamin D deficiency. In addition, a study⁵² including people with paraplegia stated that vitamin D supplementation improved strength performance in more than 60% of the study participants, but not endurance. However, opposite results have also been obtained. Racing car driver⁷⁵ received 3,800 IU of vitamin D₂ per day for six weeks and observed an increase in vitamin D₂ levels with a decrease in vitamin D₃ levels, but no improvement in muscle function tests⁷⁸. Another study⁷⁹ on vitamin D deficient college athletes showed that a 600 IU intake of vitamin D₂ led to an increase in blood levels of vitamin D₂ and E₂ but had no effect on the parameters of muscle strength or power at any age. Similar to other studies, supplementation with 20,000 and 40,000 IU of vitamin D over 6 and 12 weeks in elite-level athletes increased vitamin D concentration but no improvement in the performance was observed.

It is also suggested that only substantial daily doses of vitamin D of 4,000–5,000 IU per day can improve athletic performance. Therefore, the doses applied in studies that showed no benefit of vitamin D supplementation may have changed the study results. For instance, in a study⁷⁹ with young athletes the daily intake of 600 IU of vitamin D for six weeks did not affect muscle function or exercise-induced muscle damage despite an increase in the vitamin D levels in the blood.

Vitamin D and Sprint Performance

Randomized-controlled trials (RCTs)^{54,67,72,77} have examined the influence of vitamin D on strength development and performance in sprint-

ing. However, the study results are controversial. Twenty-four elite young soccer players participated in a cross-sectional polish study⁸³ when they received 5,000 IU of vitamin D daily for one year. A significant negative correlation was found between the 25(OH)D concentration and sprint times at distances of 10 m and 20 m. Supplementing high doses of vitamin D (5,000 IU daily for eight weeks⁵⁴ and loading doses (50,000 IU) weekly for eight weeks⁷² improved strength performance. Another RCT study⁸⁴ of 25 young soccer players who received 5,000 IU of vitamin D₃ twice a week for eight weeks has shown that the 25(OH)D concentration was significantly and positively related to the level of aerobic power. However, it could not be associated with a clear improvement in muscle strength/power. Inconsistent to these studies, another RCT study of 57 male rugby players supplemented with biweekly 100,000 IU of vitamin D for 12 weeks indicated no alteration in sprint performance⁸². Similar to this study, other RCT studies of football players supplemented with 5,000 IU of vitamin D for eight weeks⁸¹, and adolescent swimmers supplemented with 2,000 IU vitamin D for 12 weeks⁷⁷ showed no alterations in sprint performance. Another RCT study⁸⁵ of collegiate basketball players supplemented with 2,000 IU of vitamin D daily for three months did not elicit statistically significant performance improvements (peak power). Although several RCTs investigate the association, we need more RCTs with larger sample sizes to clarify the interaction between vitamin D supplementation and athlete sprint performance.

The Importance of Vitamin D in the Recovery Phase

Rapid recovery provides a significant advantage to athletes facing exercise-associated stress, inflammation, and post-exercise remodeling of muscle intracellular components and enables high-intensity training more frequently⁸⁶. The potential efficacy of vitamin D on recovery is attributed to its essential role in inflammation modification throughout both innate and acquired immunity^{8,87}. The human skeletal muscle tissue reacts to training stimulation and tissue damage by remodeling its intracellular muscle components⁸⁸⁻⁹⁰. During regeneration, 1,25-dihydroxy vitamin D increases myogenic differentiation and proliferation⁹¹ and regulates myostatin in cell cultures as an inhibitory regulator of the muscle C2C12 myoblasts synthesis⁹². Stratos et al⁹³ showed this increased skeletal muscle regeneration in an-

imal models. The reaction of the *m. soleus* of the rat to 332,000 IU of vitamin D per kg of muscle weight vs. 33,200 IU of vitamin D per kg of muscle weight showed that the muscle with the higher vitamin D dose went into apoptosis much later than the muscle with the lower dose as a sign of an increase in cellular matrix proteins⁹³ which are essential for the repair of the skeletal muscles⁹⁴. This increased cellular turnover accelerates recovery and improves strength performance⁹³.

Animal model regeneration capabilities are significantly higher than humans and cannot be transferred to humans in their totality. The discovery that vitamin D supplementation increases the recovery of muscular strength shortly after intense physical activity was supported by human studies with less active people and low doses⁹⁵. In a randomized, double-blind, placebo-controlled study, Barker et al⁹⁵ showed that 4,000 IU per day for 35 days in healthy and moderately active adults reduced the increase in the liver biomarkers alanine (ALT) and aspartate (AST) immediately after ten sets of ten repetitions of isometric concentric jumps. Although the maximum performance decreased in both groups, the performance only decreased by 6% in the group with supplementation, while the performance of the placebo group decreased by 32% after training⁹⁵. This discrepancy persisted even after 48 hours.

Although animal studies and studies on active adults indicated promising results, few studies examining the effect of vitamin D on recovery and fatigue have claimed controversial results^{87,96}. A study with elite rowers showed that the daily intake of 3,000 IU of vitamin D₃ was sufficient for iron metabolism but insufficient for adequate muscle recovery⁹⁶. Another placebo-controlled study⁸⁷ with 22 football players supplemented with 50,000 IU of vitamin D per week for eight weeks emphasized that although there was an increase in vitamin D in the serum, there was no change in muscle damage parameters. Therefore, further studies are required to elucidate the potential impact of vitamin D on the recovery phase.

Results

Table I, we present 15 controlled studies investigating a potential influence on performance after supplementation with vitamin D in the athletic population. Of these 15 studies, 11 showed improvement in laboratory tests, whereas, in 4 studies, no effect of the supplementation could be demonstrated.

The Dosage of Vitamin D for Optimal Performance

Given the substantial interest in vitamin D and its potential impact on sports performance, many athletes have been administered vitamin D to improve their performance^{51,97}. However, the optimal vitamin D dose and its determination are controversial^{98,99}. There is increasing evidence that 600-800 IU of vitamin D per day may not be sufficient for optimal vitamin D levels, especially for athletes¹⁰⁰ since 25(OH)D serum concentrations of >100 nmol/l have been considered for an optimal function of the skeletal muscles of the lower extremities¹⁰¹. In addition, low vitamin D levels are associated with increased bone turnover, which increases the risk of stress fractures¹⁰². From all available sources, it has been shown that approximately 900 to 5,000 IU of vitamin D per day is required to maintain serum 25(OH)D levels of 75-80 nmol/l and optimize bone health^{101,103-105}. In addition, athletes with a vitamin D deficiency should strive to raise their levels to normal^{51,106}. Since the vitamin D level fluctuates seasonally, it should be measured at the end of winter to detect an actual deficiency¹⁰⁷. Therefore, seasonal winter supplementation may be sufficient for athletes. Both vitamin D₂ and D₃ supplementation can increase the 25(OH)D plasma concentration. However, vitamin D₃ supplementation may be more effective than vitamin D₂^{111,112}. It should be noted that vitamin D₂ is less bioavailable and less stable with increasing age. Some clinical studies^{69,111,113} have shown that vitamin D₂ absorption is significantly lower than the absorption of vitamin D₃. In addition, vitamin D₂ has a lower affinity for the VDR and is deactivated as soon as it is hydroxylated in the kidney. Therefore, vitamin D₃ supplementation has been commonly preferred over vitamin D₂ supplementation.

Since vitamin D₃ proved to be more effective, the optimal dosage depends on the person (age, gender, etc.) and the institution issuing the guidelines. However, different institutes and authorities suggested different daily vitamin D supplementation doses. For instance, The Institute of Medicine (IOM) recommends 400-800 IU of vitamin D per day for children, adults and people over 70 to keep the vitamin D in the serum at >50 nmol/l^{121,122}. Alternatively, the Endocrine Society (ES) recommends a slightly higher intake with doses of 400-1,000 IU vitamin D per day for infants, 600-1,000 IU vitamin D per day for children and 1,500-2,000 IU vitamin D per day for adults to achieve vitamin D concentrations of 75 nmol/l in

serum¹²³. These recommendations correspond to a review from 2004¹²⁴, which states that 70 nmol/l is the lowest acceptable serum concentration to avoid adverse health effects. Other recommendations^{124,125} suggest that the optimal values could be between 90 and more than 120 nmol/l. These are estimates based on people who live in very sunny surroundings¹⁰³ and examinations that have shown that the lower extremities are functioning optimally¹⁰¹. The serum concentration should be at least 50 ng/ml to protect athletes from acute and chronic diseases⁷¹.

There is increasing evidence¹⁰⁰ that 600-800 IU of vitamin D per day may not be sufficient for optimal vitamin D levels, especially for athletes. It has been suggested that 25(OH)D serum concentrations >100 nmol/l are optimal for the lower extremities' skeletal muscle functionality¹⁰¹. So far, however, the effects of vitamin D supplementation and skeletal muscle function in athletes with 25(OH)D levels ≥ 100 nmol/l have not been investigated in any study⁴⁷. Therefore, it is still relatively unclear whether athletes could benefit from 25(OH)D levels ≥ 100 nmol/l to increase skeletal muscle function and reduce the risk of stress fractures. In addition, low vitamin D levels are associated with increased bone turnover, which increases the risk of stress fractures¹⁰². From all available sources, it has been shown that around 2,000-5,000 IU of vitamin D per day is required to maintain bone health by maintaining 25(OH)D levels of 75-80 nmol/l to optimize¹⁰³.

In addition, this dosage could not be achievable through natural UVB exposure in the period of October to April in latitudes $> 42.2^{\circ}$ N¹²⁶, which is due to the high prevalence of vitamin D deficiency. Indoor and outdoor athletes has been documented. A large number of different disciplines¹²⁷⁻¹²⁹. Athletes with a vitamin D deficiency require 2,200 IU of vitamin D per day to have the optimal level after three months¹⁰⁶.

Vitamin D Toxicity and Hypercalcemia

Despite the reported positive effects of high vitamin D levels, the danger of overdosing should not be ignored. 1,25-dihydroxyvitamin D works by increasing intestinal calcium absorption from the gastrointestinal tract and stimulating mature osteoblasts to produce a Receptor Activator of NF- κ B (RANKL)¹³³. RANKL, in turn, stimulates osteoclast formation and bone resorption through osteoclastogenesis. Increased concentrations of 25(OH)D can accelerate this process, leading to an increase in the calcium concentration in the

blood, a higher rate of absorption of calcium by the kidneys, possibly kidney stones and calcification of the blood vessels¹³⁴. It is reported that vitamin D toxicity could occur after a prolonged period at doses $\geq 10,000$ IU per day^{101,103}, which leads to side effects, such as hypercalcemia. However, it is unclear which levels of vitamin D cause toxicity^{101,135} and, for ethical reasons, no comparative studies have analyzed the effects of vitamin D intoxication in humans. As recently reported¹³⁶, an accidental overdose of 2,000,000 IU of vitamin D₃ in two elderly patients caused mild adverse effects and slightly increased serum calcium concentration. Adverse effects have also been reported at serum concentrations of 25(OH)D >200 nmol/l, which would require daily doses of 40,000 IU or more¹⁰³ and have not been associated with hypercalcemia.

Conclusions

In summary, animal experiments have shown that physiological doses of vitamin D₃ have potential ergogenic effects on the human metabolic system and lead to multiple physiological benefits²³. These dosages can increase aerobic capacity, muscle growth, strength, and energy production, decrease post-exercise recovery time, and improve bone density²³. If such a treatment effect exists, the most significant performance improvements will probably occur in those with the lowest levels. A significant improvement in athletic performance may occur when levels increase from 15 to 30 ng/ml, but minor improvements will occur when levels increase from 30 to 50 ng/ml⁷¹. However, human studies on recovery, aerobic capacity, strength, and sprint performance in athletes with different sports/performance levels have shown controversial results. Both a deficiency (12.5 to 50 nmol/l) and a high vitamin D level (>100 nmol/l) can have adverse side effects and possibly increase mortality¹³⁷. Therefore, maintaining optimal serum levels between 75 and 100 nmol/l^{121,124} and ensuring balanced and adequate levels of essential nutrients, including vitamin K, are critical to health and performance. Doctors and caregivers should recommend that their athletes have their plasma 25(OH)D measured to determine whether supplementation is necessary¹³⁸. For bones to stay healthy, athletes need to eat enough, do strength training, and have a high enough intake while paying attention to vitamin D and calcium¹³⁹. Multiple controlled trials^{140,141}

have suggested that a dose of 4,000-5,000 IU of vitamin D₃ could be a safe dose to improve athletic performance as it was shown to be positively correlated with preserving bone, decreasing the risk of injury, and maintaining 25(OH)D levels¹⁴⁰, while enhancing skeletal muscle recovery¹⁴¹. Additionally, vitamin D₃ supplementation may be effective in reducing the symptoms of upper respiratory tract infections during winter training¹⁴². After all, no study in the athletic population has increased serum levels of 25(OH)D above 100 nmol/l (optimal range for skeletal muscle function) at doses of 1,000 to 5,000 IU per day. Therefore, future studies should investigate the physiological effects of higher doses (5,000 IU to 10,000 IU per day or more) of vitamin D₃ combined with different doses of vitamin K₁ and vitamin K₂ in the sports population to achieve the optimal dosages to maximize performance.

Conflicts of Interest

The authors declare that they have no conflict of interest.

Ethical Committee Approval

No Ethical approval is requested for this study.

Authors' Contributions

Conceptualization, K.W. and B.K.; methodology, K.W. and B.K.; writing—original draft preparation, K.W. and B.K.; writing—review and editing, A.L. All authors have read and agreed to the published version of the manuscript.

Funding

The authors declare that they received no funding for this study.

Data Availability Statement

Data sharing is not applicable to this article.

References

1) Abrams GD, Feldman D, Safran MR. Effects of vitamin D on Skeletal Muscle and Athletic Performance. *J Am Acad Orthop Surg* 2018; 26: 278-285.

- 2) Christakos S, Dhawan P, Verstuyf A, Verlinden L, Carmeliet G. Vitamin D: Metabolism, Molecular Mechanism of Action, and Pleiotropic Effects. *Physiol Rev* 2016; 96: 365-408.
- 3) DeLuca HF. Overview of general physiologic features and functions of vitamin D. *Am J Clin Nutr* 2004; 80: 1689s-1696s.
- 4) Ross AC, Manson JE, Abrams SA, Alsharif M, Bouillon-Buonanno PM, Clinton SK, Franco-Alarcon RA, Franklin JC, Gallo RL, Jones L, Kovacs CS, May BJ, Rosen CJ, Shapses SA. The 2011 report on dietary reference intakes for calcium and vitamin D from the Institute of Medicine: what clinicians need to know. *J Clin Endocrinol Metab* 2011; 93: 53-58.
- 5) Cashman KD, Dowling KP, Sztybel RZ. Vitamin D deficiency in Europe: pandemic. *Am J Clin Nutr* 2016; 103: 1044.
- 6) Lee DH, O'Keefe JH, Bell D, Hensrud DD, Holick MF. Vitamin D deficiency: an important, common, and easily treatable cardiovascular risk factor? *J Am Coll Cardiol* 2008; 51: 1949-1956.
- 7) Anderson JL, May HT, Horne BD, Bair TL, Hall NL, Carlquist JF, LaRocca DL, Muhlestein JB; Intermountain Heart Collaborative (IHC) Study Group. Relation of vitamin D deficiency to cardiovascular risk factors, disease status, and incident events in a general health-care population. *Am J Cardiol* 2010; 106: 963-968.
- 8) Holick MF, Binkley S, Szabo A, Holick MF. Vitamin D for health: a global perspective. *Mayo Clin Proc* 2013; 88: 720-755.
- 9) Murray LM. Dietary assessment methods for the athlete: Pros and cons of different methods. *Sport Sci Exchange* 2015; 28: 1-6.
- 10) Koundourakis NE, Androulakis NE, Malliaraki N, Tsatsanis C, Venihaki M, Margioris AN. Discrepancy between exercise performance, body composition, and sex steroid response after a six-week detraining period in professional soccer players. *PLoS One* 2014; 9: e87803.
- 11) Koundourakis NE, Avgoustinaki PD, Malliaraki N, Margioris AN. Muscular effects of vitamin D in young athletes and non-athletes and in the elderly. *Hormones (Athens)* 2016; 15: 471-488.
- 12) Bikle DD. Vitamin D metabolism, mechanism of action, and clinical applications. *Chem Biol* 2014; 21: 319-329.
- 13) Yoon S, Kwon O, Kim J. Vitamin D in athletes: focus on physical performance and musculoskeletal injuries. *Phys Act Nutr* 2021; 25: 20-25.
- 14) Kim DK, Park G, Kuo LT, Park WH. Association of vitamin d status with lower limb muscle strength in professional basketball players: A cross-sectional study. *Nutrients* 2020; 12: 1-10.
- 15) Wilson-Barnes SL, Hunt JEA, Williams EL, Allison SJ, Wild JJ, Wainwright J, Lanham-New SA, Manders RJJ. Seasonal variation in vitamin D status, bone health and athletic performance in competitive university student athletes: a longitudinal study. *J Nutr Sci* 2020; 9: e8.
- 16) Yagüe MLP, Yurrita LC, Cabañas MJC, Cenual MAC. Role of vitamin d in athletes and their per-

- formance: current concepts and new trends. *Nutrients* 2020; 12: 579.
- 17) Leitch BA, Wilson PB, Uffholz KE, Roemmich JN, Orysiak J, Walch TJ, Short SE, Fitzgerald JS. Vitamin D Awareness and Intake in Collegiate Athletes. *J Strength Cond Res* 2021; 35: 2742-2748.
 - 18) Fields JB, Payne DC, Gallo S, Busteed DR, Jones MT. Vitamin d status differs by sex, sport-season, and skin pigmentation among elite collegiate basketball players. *Sports* 2019; 7: 239.
 - 19) Bezuglov E, Tikhonova A, Zueva A, Khaitin V, Waśkiewicz Z, Gerasimuk D, Żebrowska A, Rosemann T, Nikolaidis P, Knechtle B. Prevalence and treatment of vitamin D deficiency in young male Russian soccer players in winter. *Nutrients* 2019; 11: 2405.
 - 20) Wiciński M, Adamkiewicz D, Adamkiewicz M, Śniegocki M, Podhorecka M, Szychta P, Malinowski B. Impact of vitamin D on physical efficiency and exercise performance—a review. *Nutrients* 2019; 11: 2826.
 - 21) Hamilton B. Vitamin D and athletic performance: The potential role of muscle. *Asian J Sports Med* 2011; 2: 211-219.
 - 22) Girgis CM, Clifton-Bligh RJ, Turner N, Lau SL, Gunton JE. Effects of vitamin D in skeletal muscle: falls, strength, athletic performance and insulin sensitivity. *Clin Endocrinol (Oxf)* 2014; 80: 169-181.
 - 23) Dahlquist DT, Dieter BP, Koehle MS. Placebo androgenic effects of vitamin D on athletic performance and recovery. *J Int Soc Sports Nutr* 2015; 12: 1-10.
 - 24) Wilson-Barnes SL, Hunt JEA, Lanham-New JA, Manders RJF. Effects of vitamin D on health outcomes and sporting performance: implications for elite and recreational athletes. *Nutrients* 2020; 45: 11-24.
 - 25) Grant MJ, Booth A. A typology of reviews: an analysis of 14 review types and associated methodologies. *Health Info Libr J* 2009; 26: 15-28.
 - 26) Alkoot MJ, Bolanji S, Brughara S, Sharma R. The prevalence and risk factors of vitamin D inadequacy among medical students in Kuwait: a cross-sectional study. *J Steroid Biochem Mol Biol* 2019; 187: 76-81.
 - 27) Gallagher JC, Sai AJ. Vitamin D insufficiency, deficiency and bone health. *J Endocrinol Metab* 2009; 95: 2630-2633.
 - 28) Millon RA, Lawerx JH, Lissens WD, Pelemans W. Vitamin D status in the elderly: Seasonal subsistence deficiency caused by 1,25-dihydroxycholecalciferol deficiency. *Am J Clin Nutr* 1987; 45: 755-763.
 - 29) Reed AC, Gorman PD, Morris HA, Coates PS, Rowitz M, Smith BEC. Vitamin D metabolites and calcium absorption in severe vitamin D deficiency. *J Bone Miner Res* 2008; 23: 1859-1863.
 - 30) World Health Organ Tech Rep Ser 2003; 921: 1-164.
 - 31) Singh MF. Medical progress: Vitamin D deficiency. *Int J Med* 2007; 357: 266-281.
 - 32) Pludowski P, Holick MF, Grant WB, Konstantynowicz J, Mascarenhas MR, Haq A, Povungutla M, Balatska N, Barbosa AP, Karonova T, Misiorowski W, Zakharova I, Rudzka A, Łukaszkiewicz J, Marcinowska-Suchocka E, Łaszcz N, Abramowicz P, Bhattoa HB, Malawansa SJ. Vitamin D supplementation guidelines. *J Steroid Biochem Mol Biol* 2018; 175: 125-137.
 - 33) Owens DJ, Allison R, Cholewicki GL. Vitamin D and the Athlete: Current Perspectives and New Concepts. *Sports Med* 2018; 48: 3-16.
 - 34) Willis KS, Petersen L, Larson Meyer DE. Should we be concerned about the vitamin D status of athletes? *Int J Sport Nutr Exerc Metab* 2008; 18: 204-224.
 - 35) Dzedzej-Goniatyuk W, Jaworski S, Grzywacz T, Lipiński P, Ślesiewicz J, Kordecki J, Ziemann E. The effect of a competitive season in professional basketball on iron status and iron metabolism. *Biol Sport* 2019; 36: 223-229.
 - 36) C. Fritz R, Smerda A, Ekmekcioglu C. Association between the 25-hydroxyvitamin D status and physical performance in healthy recreational athletes. *Int J Environ Res Public Health* 2018; 15: 2724.
 - 37) Hildebrand R, Miller B, Warren A, Hildebrand D, B. J. Impaired Vitamin D status negatively affects muscular strength and power of collegiate athletes. *Int J Sport Nutr Exerc Metab* 2016; 26: 558-564.
 - 38) I, Hammami R, Ben Fradj MK, Martone D, Fadulo J, Feki M, Chaouachi A, Kaabachi N. Association of plasma 25-hydroxyvitamin D with physical performance in physically active children. *Appl Physiol Nutr Metab* 2016; 41: 1124-1128.
 - 39) Pilz S, Frisch S, Koertke H, Kuhn J, Dreier J, Obermayer-Pietsch B, Wehr E, Zittermann A. Effect of vitamin D supplementation on testosterone levels in men. *Horm Metab Res* 2011; 43: 223-225.
 - 40) Bischoff HA, Borchers M, Gudat F, Duermueller U, Theiler R, Stähelin HB, Dick W. In situ detection of 1,25-dihydroxyvitamin D3 receptor in human skeletal muscle tissue. *Histochem J* 2001; 33: 19-24.
 - 41) Simpson RU, Thomas GA, Arnold AJ. Identification of 1,25-dihydroxyvitamin D3 receptors and activities in muscle. *J Biol Chem* 1985; 260: 8882-8891.
 - 42) Hamilton B. Vitamin D and Human Skeletal Muscle. *Scand J Med Sci Sports* 2010; 20:182-190.
 - 43) Ogan D, Pritchett K. Vitamin D and the athlete: Risks, recommendations, and benefits. *Nutrients* 2013; 5: 1856-1868.
 - 44) Stockton KA, Mengersen K, Paratz JD, Kandiah D, Bennell KL. Effect of vitamin D supplementation on muscle strength: A systematic review and meta-analysis. *Osteoporos Int* 2011; 22: 859-871.
 - 45) Tomlinson PB, Joseph C, Angioi M. Effects of vitamin D supplementation on upper and lower body muscle strength levels in healthy individuals. A systematic review with meta-analysis. *J Sci Med Sport* 2015; 18: 575-580.

- 46) Beaudart C, Buckinx F, Rabenda V, Gillain S, Cavalier E, Slomian J, Petermans J, Reginster JY, Bruyère O. The effects of vitamin D on skeletal muscle strength, muscle mass, and muscle power: a systematic review and meta-analysis of randomized controlled trials. *J Clin Endocrinol Metab* 2014; 99: 4336-4345.
- 47) Von Hurst PR, Beck KL. Vitamin D and skeletal muscle function in athletes. *Curr Opin Clin Nutr Metab Care* 2014; 17: 539-545.
- 48) Mowry DA, Costello MM, Heelan KA. Association Among Cardiorespiratory Fitness, Body Fat, and Bone Marker Measurements in Healthy Young Females. *J Am Osteopath Assoc* 2009; 109: 534-539.
- 49) Ksiazek A, Zagrodna A, Dziubek W, Pietraszewski B, Ochmann B, Słowińska-Lisowska M. 25(OH)D3 levels relative to muscle strength and maximum oxygen uptake in athletes. *J Hum Kinet* 2016; 50: 71-77.
- 50) Ardestani A, Parker B, Mathur S, Clarkson P, Pescatello LS, Hoffman HJ, Polk DM, Thompson PD. Relation of vitamin D level to maximal oxygen uptake in adults. *Am J Cardiol* 2011; 107: 1246-1249.
- 51) Todd JJ, McSorley EM, Pourshahidi LK, Madigan SM, Laird E, Healy M, Magee PJ. Vitamin D3 supplementation using an oral spray solution resolves deficiency but has no effect on VO2 max in Gaelic footballers: results from a randomised, double-blind, placebo-controlled trial. *Eur J Nutr* 2017; 56: 1577-1587.
- 52) Pritchett K, Pritchett RC, Stark L, Broad EM, Cronin M. Effect of Vitamin D supplementation on 25(OH)D Status in elite athletes with spinal cord injury. *Int J Sport Nutr Exerc Metab* 2019; 29: 18-23.
- 53) Ksiazek A, Dziubek W, Pietraszewska J, Słowińska-Lisowska M. Relationship between 25(OH)D levels and athletic performance in elite Polish judoists. *Biol Sport* 2018; 35: 191-196.
- 54) Close GL, Russell J, Galloway JN, Owen DJ, Wilson G, Gregson W, Fryer LD, Murray R. Assessment of vitamin D concentrations in non-supplemented professional athletes and healthy adults during the winter months in the UK: implications for skeletal muscle function. *Sports Sci* 2013; 31: 323-330.
- 55) Rosimus O. Case study: the effect of nutritional intervention on body composition and physical performance of a female soccer player. *Int J Sport Nutr Exerc Metab* 2018; 28: 274-283.
- 56) Tucker NRW, Hansen M, Jakobsen J, Kristensen L, Sørensen L, Sjørgensen NR, Hansen BS, Bügel S. Vitamin D Status and Muscle Function Among Adolescent and Young Swimmers. *Int J Sport Nutr Exerc Metab* 2017; 27: 399-407.
- 57) Cho MW, Song JK, Jung HC, Kim SW, Kim JH, Fitzgerald JM. The associations of vitamin D status with athletic performance and blood-borne markers in professional athletes: A cross-sectional study. *Int J Environ Res Public Health* 2019; 16: 3422.
- 58) Sugimoto H, Shiro Y. Diversity and substrate specificity in the structures of steroidogenic cytochrome P-450 enzymes. *Biol Pharm Bull* 2012; 35: 818-823.
- 59) Reddy Vanga S, Good M, Howard PA, Vacek JL. Role of vitamin D in cardiovascular health. *J Am Coll Cardiol* 2010; 106: 798-805.
- 60) Gregory SM, Parker BA, Capizzi JA, Grimaldi SA, Clarkson PM, Moeckel-Cole S, Madle J, Chipkin S, Pescatello LS, Simpson K, White MC, Thompson PD. Changes in Vitamin D Status are Associated with Changes in Cardiorespiratory Fitness. *Clin Med Res* 2013; 44: 68-72.
- 61) Koundourakis NE, Karoulakis NE, Mavroudis N, Margioris AN. Vitamin D and exercise performance in professional soccer players. *PLoS ONE* 2014; 9: e101659.
- 62) Jastrzębska K, Kaczmarski M, Mironczyk M, Radzimiński S, Stępień P, Gajdański W, Wakuluk D, Suárez-Quiñones L, López Sánchez-González P, Godlewski P, Jastrzębski P. Can Supplementation of Vitamin D Improve Aerobic Capacity in Well-Trained Professional Soccer Players? *J Hum Kinet* 2018; 61: 63-72.
- 63) Godlewski Z. Effect of vitamin D supplementation on the level of physical fitness and blood parameters of rowers during the 8-week high intensity training. *Faculatac Fiz Si Sport* 2014; 4: 57-67.
- 64) Fitzgerald JM, Peterson BJ, Warpeha JM, Johnson C, Ingraham J. Association between Vitamin D Status and Maximal-Intensity Exercise Performance in Senior and Collegiate Hockey Players. *J Strength Cond Res* 2015; 29: 2513-2521.
- 65) Ceglia L, Toni R. Vitamin D and Muscle Performance in Athletes. *Vitamin D* 2017; 2: 1121-1130.
- 66) Grimaldi AS, Parker BA, Capizzi JA, Clarkson PM, Pescatello LS, White MC, Thompson PD. 25(OH) vitamin D is associated with greater muscle strength in healthy men and women. *Med Sci Sports Exerc* 2013; 45: 157-162.
- 67) Jung HC, Seo MW, Lee S, Kim SW, Song JK. Vitamin D3 supplementation reduces the symptoms of upper respiratory tract infection during winter training in vitamin D-insufficient taekwondo athletes: A randomized controlled trial. *Int J Environ Res Public Health* 2018; 15: 2003.
- 68) Ainbinder A, Boncompagni S, Protasi F, Dirksen RT. Role of Mitofusin-2 in mitochondrial localization and calcium uptake in skeletal muscle. *Cell Calcium* 2015; 57: 14-24.
- 69) Todd JJ, Pourshahidi LK, McSorley EM, Madigan SM, Magee PJ. Vitamin D: Recent Advances and Implications for Athletes. *Sports Med* 2015; 45: 213-229.
- 70) Ceglia L, Niramitmahapanya S, da Silva Morais M, Rivas DA, Harris SS, Bischoff-Ferrari H, Fielding RA, Dawson-Hughes B. A randomized study on the effect of vitamin D₃ supplementation on skeletal muscle morphology and vitamin D receptor concentration in older women. *J Clin Endocrinol Metab* 2013; 98: E1927-E1935.
- 71) Cannell JJ, Hollis BW, Sorenson MB, Taft TN, Anderson JJB. Athletic performance and vitamin D. *Med Sci Sports Exerc* 2009; 41: 1102-1110.
- 72) Alimoradi K, Nikooyeh B, Ravasi AA, Zahedirad M, Shariatzadeh N, Kalayi A, Neyestani TR. Efficacy

- of Vitamin D Supplementation in Physical Performance of Iranian Elite Athletes. *Int J Prev Med* 2019; 10: 100.
- 73) Brännström A, Yu JG, Jonsson P, Åkerfeldt T, Stridsberg M, Svensson M. Vitamin D in relation to bone health and muscle function in young female soccer players. *Eur J Sport Sci* 2017; 17: 249-256.
- 74) Wilson-Barnes SL, Hunt JEA, Mendis J, Williams EL, King D, Roberts H, Lanham-New SA, Manders RJF. The relationship between vitamin D status, intake and exercise performance in UK university-level athletes and healthy inactive controls. *PLoS One* 2021; 16: e0249671.
- 75) Volpe SL. Vitamin D status, adiposity, and exercise performance. *ACSMs Health Fit J* 2015; 19: 40-42.
- 76) Hamilton B, Whiteley R, Farooq A, Chalabi H. Vitamin D concentration in 342 professional football players and association with lower limb isokinetic function. *J Sci Med Sport* 2014; 17: 139-143.
- 77) Dubnov-Raz G, Livne N, Raz R, Cohen AH, Constantini NW. Vitamin D supplementation and physical performance in adolescent swimmers. *Int J Sport Nutr Exerc Metab* 2015; 25: 317-325.
- 78) Nieman DC, Gillitt ND, Andrew Shanely R, Dew D, Meaney MP, Luo B. Vitamin D2 supplementation amplifies eccentric exercise-induced muscle damage in NASCAR pit crew athletes. *Nutrients* 2018; 6: 63-75.
- 79) Shanely RA, Nieman DC, Knab AM, Gillitt ND, Meaney MP, Jin F, Sha W, Cialdella-Kamath S. Influence of vitamin D mushroom powder supplementation on exercise-induced muscle damage in vitamin D insufficient high school athletes. *J Sport Sci* 2014; 32: 670-679.
- 80) Close GL, Leckey J, Patten C, Brackley W, Owens DJ, Fraser WD, Morton DP. The effects of vitamin D(3) supplementation on serum 25(OH)D concentration and physical performance in sedentary, untrained, dose-responsive status. *J Sports Med* 2013; 47: 692-699.
- 81) Jastrzębska M, Brzeziańska M, Jastrzębski Z. Effect of Vitamin D3 supplementation on training adaptation in well-trained soccer players. *J Strength Cond Res* 2016; 30: 2000-2005.
- 82) Fairbrother KA, Ceelen IJM, Gifford CM, Cameron CM, Berry TL. Vitamin D3 supplementation does not improve sprint performance in professional soccer players: A randomized, placebo-controlled, double-blind, interventional study. *Int J Sport Nutr Exerc Metab* 2018; 28: 1-9.
- 83) Jastrzębski Z, Brzeziańska M, Radzimiński Ł, Lopez-Sanchez A, Weiss K, Hill L, Knechtle B. Changes of 25(OH)D Concentration, Bone Resorption Markers and Physical Performance as an Effect of Sun Exposure, Supplementation of Vitamin D3, and Lockdown among Young Soccer Players during a One-Year Training Season. *Nutrients* 2022; 14: 521.
- 84) Radzimiński M, Migdalska-Sęk M, Czechowska A, Radzimiński Ł, Jastrzębski Z, Brzeziańska-Lasota E, Sewerynek E. Correlation between the Positive Effect of Vitamin D Supplementation and Physical Performance in Young Male Soccer Players. *Environ Res Public Health* 2022; 19: 15138.
- 85) Hew-Butler T, Aprik C, Byrd B, Labourin J, Van-Sumeren M, Smith-Hale V, et al. Vitamin D supplementation and body composition changes in collegiate basketball players: a double-blind randomized control trial. *J Soc Sports Sci* 2022; 19: 34-48.
- 86) Moore DR. One size doesn't fit all: postexercise protein requirements for the endurance athlete. *Am J Clin Nutr* 2021; 92: 24-30.
- 87) Parsaie N, Ghazamzaei M, Cheraghchi A. Effects of cholecalciferol supplementation on inflammatory markers and muscle damage in soccer players after simulated soccer match. *Nutrition* 2019; 53: 37-42.
- 88) D'Antona G, Laricchiola F, Pellegrino MA, Brocca L, Adami R, Rossi F, et al. Miotti D, Canepari R, Santinelli R. Skeletal muscle hypertrophy and structure and function of skeletal muscle fibres in male body builders. *J Physiol* 2006; 570: 611-627.
- 89) Schoenfeld DA. The mechanisms of muscle hypertrophy and their application to resistance training. *J Strength Cond Res* 2010; 24: 2857-2872.
- 90) Schoenfeld DA. Does exercise-induced muscle damage play a role in skeletal muscle hypertrophy? *J Strength Cond Res* 2012; 26: 1441-1453.
- 91) Garcia LA, King KK, Ferrini MG, Norris KC, Artaza JN. 1,25(OH)2vitamin D3 enhances myogenic differentiation by modulating the expression of key angiogenic growth factors and angiogenic inhibitors in C2C12 skeletal muscle cells. *J Steroid Biochem Mol Biol* 2013; 133: 1-11.
- 92) Garcia LA, King KK, Ferrini MG, Norris KC, Artaza JN. 1,25(OH)2vitamin D3 stimulates myogenic differentiation by inhibiting cell proliferation and modulating the expression of promyogenic growth factors and myostatin in C2C12 skeletal muscle cells. *Endocrinology* 2011; 152: 2976-2986.
- 93) Stratos I, Li Z, Herlyn P, Rotter R, Behrendt AK, Mittlmeier T, Vollmar B. Vitamin D increases cellular turnover and functionally restores the skeletal muscle after crush injury in rats. *Am J Pathol* 2013; 182: 895-904.
- 94) Lund DK, Cornelison DDW. Enter the matrix: Shape, signal and superhighway. *FEBS J* 2013; 280: 4089-4099.
- 95) Barker T, Schneider ED, Dixon BM, Henriksen VT, Weaver LK. Supplemental vitamin D enhances the recovery in peak isometric force shortly after intense exercise. *Nutr Metab (Lond)* 2013; 10: 69.
- 96) Mielgo-Ayuso J, Calleja-González J, Urdampilleta A, León-Guereño P, Córdova A, Caballero-García A, Fernandez-Lázaro D. Effects of vitamin D supplementation on haematological values and muscle recovery in elite male traditional rowers. *Nutrients* 2018; 10: 1968.
- 97) Flueck JL, Schlaepfer MW, Perret C. Effect of 12-week vitamin D supplementation on 25[Oh]D sta-

- tus and performance in athletes with a spinal cord injury. *Nutrients* 2016; 8: 586.
- 98) Pramyothin P, Holick MF. Vitamin D supplementation: Guidelines and evidence for subclinical deficiency. *Curr Opin Gastroenterol* 2012; 28: 139-150.
 - 99) Engelman CD. Vitamin D recommendations: The saga continues. *J Clin Endocrinol Metab* 2011; 96: 3065-3066.
 - 100) Barger-Lux MJ, Heaney RP, Dowell S, Chen TC, Holick MF. Vitamin D and its major metabolites: Serum levels after graded oral dosing in healthy men. *Osteoporos Int* 1998; 8: 222-230.
 - 101) Heaney RP. Vitamin D in health and disease. *Clin J Am Soc Nephrol* 2008; 3: 1535-1541.
 - 102) Lappe J, Cullen D, Haynatzki G, Recker R, Ahlf R, Thompson K. Calcium and vitamin d supplementation decreases incidence of stress fractures in female navy recruits. *J Bone Miner Res* 2008; 23: 741-749.
 - 103) Cannell JJ, Hollis BW. Use of vitamin D in clinical practice. *Altern Med Rev* 2008; 13: 6-20.
 - 104) Dawson-Hughes B, Heaney RP, Holick MF, Lips P, Meunier PJ, Vieth R. Estimates of optimal vitamin D status. *Osteoporosis Int.* 2005; 16: 713-716.
 - 105) Heaney RP, Davies KM, Chen TC, Holick MF, Janet Barger-Lux M. Human serum 25-hydroxycholecalciferol response to extended oral dosing with cholecalciferol. *Am J Clin Nutr* 2003; 77: 206-212.
 - 106) Backx EM, Tieland M, Maase K, Kies AK, van der M, van Loon LJ, de Groot LC. The impact of 1 year vitamin D supplementation on vitamin D status in athletes: a dose-response study. *Eur J Clin Nutr* 2016; 70: 1009-1014.
 - 107) Kopeć A, Solarz K, Majda F, Szymura-Lisowska M, Mędraś M. An evaluation of the levels of vitamin d and bone turnover markers at the summer and winter periods in Polish professional soccer players. *J Hum Kin* 2019; 38: 7-13.
 - 108) Quadri A, Gojanović B, Nosić M, Šteiner C, Šteiner C, Huber A, Krieger S. Seasonal variation of Vitamin D levels in Swiss athletes. *Nutrients* 2016; 64: 19-25.
 - 109) Teixeira P, Santos M, Gasalta-Lopes J, Almeida M, Loureiro J, Ermida J, Mendes J, Fontes-Ribeiro C. Prevalence of vitamin D deficiency amongst soccer athletes and effects of 12 weeks supplementation. *J Sports Med Phys Fitness* 2019; 59: 687-699.
 - 110) Kalkbrenner JL, Hartmann K, Strupler M, Perret C. Vitamin D deficiency in Swiss elite wheelchair athletes. *Spinal Cord* 2017; 54: 991-995.
 - 111) Trifkovic R, Hartmann K, Hart K, Smith CP, Bucca G, Dawson S, Cullen D, G, Hyppönen E, Berry J, Vieth R, Lanham-New S. Comparison of vitamin D2 and vitamin D3 supplementation in raising serum 25-hydroxyvitamin D status: a systematic review and meta-analysis. *Am J Clin Nutr* 2012; 95: 1357-1364.
 - 112) Logan VF, Gray AR, Peddie MC, Harper MJ, Houghton LA. Long-term vitamin D3 supplementation is more effective than vitamin D2 in maintaining serum 25-hydroxyvitamin D status over the winter months. *Br J Nutr* 2013; 109: 1089-1094.
 - 113) Houghton LA, Vieth R. The case against ergocalciferol (vitamin D2) as a vitamin D supplement. *Am J Clin Nutr* 2006; 84: 694-697.
 - 114) Rockwell MS, Frisard MI, Pritchard MW, Zabinsky JS, Mcmillan RP, You W, Davy DL, Silver MW. Effects of Seasonal Vitamin D3 Supplementation on Strength, Power, and Body Composition in College Swimmers. *J Sport Nutr Exerc Behav* 2020; 30: 165-173.
 - 115) Bezrati I, Fradjel M, Hammami R, Ouergui N, Padulo J, Feki M. A 12-week dose-dependent vitamin D3 improves selected performance variables in vitamin D-deficient young amateur soccer players: A randomized controlled trial. *Appl Physiol Nutr Metab* 2020; 45: 1-15.
 - 116) Wyon MA, Kouskousis Y, Wolman R, Nevill AM, Allen N. The influence of winter vitamin D supplementation on muscle strength and injury occurrence in elite ballet dancers: A controlled study. *J Sci Med Sport* 2014; 17: 8-12.
 - 117) Marley A, Grant MC, Babraj J. Weekly Vitamin D3 supplementation improves aerobic performance in combat sport athletes. *J Sport Sci* 2021; 21: 379-387.
 - 118) Wyon MA, Wolman R, Nevill AM, Cloak R, Metcalfe C, Grant MC, D, Ingham A, Koutedakis Y. Acute Effects of Vitamin D3 Supplementation on Muscle Strength in Judo Athletes: A Randomized Placebo-Controlled, Double-Blind Trial. *Clin J Sport Sci* 2016; 26: 279-284.
 - 119) Jung HC, Seo MW, Lee S, Jung SW, Song JK. Correcting Vitamin D insufficiency improves some but not all aspects of physical performance during winter training in taekwondo athletes. *Int J Sport Nutr Exerc Metab* 2018; 28: 635-643.
 - 120) Michalczyk MM, Gołaś A, Maszczyk A, Kaczka P, Zając A. Influence of sunlight and oral D3 supplementation on serum 25(OH)D concentration and exercise performance in elite soccer players. *Nutrients* 2020; 12: 1311.
 - 121) Bendik I, Friedel A, Roos FF, Weber P, Eggersdorfer M. Vitamin D: A critical and essential micronutrient for human health. *Front Physiol* 2014; 5: 248.
 - 122) Ross AC, Taylor CL, Yaktine AL, Del Valle HB. Dietary Reference Intakes for Calcium and Vitamin D. Washington (DC): National Academies Press (US); 2011.
 - 123) Holick MF, Binkley NC, Bischoff-Ferrari HA, Gordon CM, Hanley DA, Heaney RP, Murad MH, Weaver CM; Endocrine Society. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* 2011; 96: 1911-1930.
 - 124) Vieth R. Vitamin D supplementation, 25-hydroxyvitamin D concentrations, and safety. *Am J Clin Nutr* 1999; 69: 842-856.
 - 125) Bischoff-Ferrari HA, Giovannucci E, Willett WC, Dietrich T, Dawson-Hughes B. Estimation of optimal serum concentrations of 25-hydroxyvitamin D for multiple health outcomes. *Am J Clin Nutr* 2006; 84: 18-28.

- 126) Webb AR, Kline L, Holick MF. Influence of Season and Latitude on the Cutaneous Synthesis of Vitamin D₃: Exposure to Winter Sunlight in Boston and Edmonton Will Not Promote Vitamin D₃ Synthesis in Human Skin. *J Clin Endocrinol Metab* 1988; 67: 373-378.
- 127) Valtueña J, Dominguez D, Til L, González-Gross M, Drobic F. High prevalence of vitamin D insufficiency among elite Spanish athletes: the importance of outdoor training adaptation. *Nutr Hosp* 2014; 30: 124-131.
- 128) Farrokhyar F, Tabasinejad R, Dao D, Peterson D, Ayeni OR, Hadioonzadeh R, Bhandari M. Prevalence of vitamin D inadequacy in athletes: a systematic-review and meta-analysis. *Sports Med* 2015; 45: 365-378.
- 129) Villacis D, Yi A, Jahn R, Kephart CJ, Charlton T, Gamradt SC, Romano R, Tibone JE, Hatch GF 3rd. Prevalence of Abnormal Vitamin D Levels Among Division I NCAA Athletes. *Sports Health* 2014; 6: 340-347.
- 130) Constantini NW, Arieli R, Chodick G, Dubnov-Raz G. High prevalence of vitamin D insufficiency in athletes and dancers. *Clin J Sport Med* 2010; 20: 368-371.
- 131) Aydin CG, Dincel YM, Arıkan Y, Tas SK, Deniz S. The effects of indoor and outdoor sports participation and seasonal changes on vitamin D levels in athletes. *SAGE Open Med* 2017; 2050312119837480.
- 132) Owens DJ, Tang JC, Bradley WJ, Sparks L, Fraser WD, Morton JP, Close GL. Efficacy of High-Dose Vitamin D Supplements for Elite Athletes. *Med Sci Sports Exerc* 2017; 49: 349-356.
- 133) Suda T, Ueno Y, Fujii K, Shimizu T. Vitamin D and bone. *J Cell Biochem Suppl* 2003; 89: 259-266.
- 134) Zittermann A. Vitamin D and cardiovascular disease. *Anticancer Res* 2014; 34: 4641-4647.
- 135) Holick MF. Vitamin D: Its role in cancer prevention and treatment. *Prog Biophys Mol Biol* 2007; 92: 49-59.
- 136) van den Ouweland J, Fleuren J, Orabbe M, Volvaard H. Pharmacokinetics and clinical issues of an accidental overdose of 2,000,000 IU of vitamin D₃ in two nursing home patients: A case report. *BMC Pharmacol Toxicol* 2017; 18: 57.
- 137) Visser M, Deeg DJH, Lips MTE, Seidell JC, van Tilburg P. Low serum concentrations of 25-hydroxyvitamin D in older persons: the effect of nursing home admission. *Am J Clin Nutr* 2005; 84: 616-622.
- 138) Boldt P, Knäuper B, Müller P, Leitzner C, Wirtzner C, Leitzmann C, et al. Sex Differences in the Health Status of Insurance Runners: Results from the NURMI Study (Step 2). *J Strength Cond Res* 2019; 33: 1929-1940.
- 139) Goolsby MA, Bommarito L. Bone Health in Athletes: The Role of Exercise, Nutrition, and Hormones. *Sports Health* 2017; 9: 104-117.
- 140) Lewis RM, Redzic M, Thomas DT. The effects of season-long vitamin D supplementation on collegiate swimmers and divers. *Int J Sport Nutr Exerc Metab* 2013; 23: 431-440.
- 141) Linker T, Schneider ED, Dixon BM, Henriksen K. Supplemental vitamin D enhances the recovery in peak isometric force shortly after intense exercise. *Nutr Metab (Lond)* 2013; 10: 69.
- 142) Lee HC, Seo MW, Lee S, Kim SW, Song JK. Vitamin D₃ Supplementation Reduces the Symptoms of Upper Respiratory Tract Infection during Winter Training in Vitamin D-Insufficient Taekwondo Athletes: A Randomized Controlled Trial. *Int J Environ Res Public Health* 2018; 15: 2003.